

Case No.: ITT0061P
 Filed: 04/06/2006
 US National Phase of
 PCT/GB2004/004306

(19) World Intellectual Property Organization
 International Bureau



(43) International Publication Date
 6 February 2003 (06.02.2003)

PCT

(10) International Publication Number
WO 03/010140 A2

(51) International Patent Classification⁷: **C07D 209/00**

[CA/CA]; 2100 Cunard Street, Laval, Québec H7S 2G5 (CA).

(21) International Application Number: **PCT/CA02/01127**

(22) International Filing Date: **18 July 2002 (18.07.2002)**

(74) Agent: **BERNIER, Louise, G.**; Boehringer Ingelheim (Canada) Ltd., 2100 Cunard Street, Laval, Québec H7S 2G5 (CA).

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
 60/307,674 25 July 2001 (25.07.2001) US
 60/338,061 7 December 2001 (07.12.2001) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*):
BOEHRINGER INGELHEIM (CANADA) LTD.
 [CA/CA]; 2100 Cunard Street, Laval, Québec H7S 2G5 (CA).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BEAULIEU, Pierre, Louis** [CA/CA]; 2100 Cunard Street, Laval, Québec H7S 2G5 (CA). **FAZAL, Gulrez** [CA/CA]; 2100 Cunard Street, Laval, Québec H7S 2G5 (CA). **KUKOLJ, George** [CA/CA]; 2100 Cunard Street, Laval, Québec H7S 2G5 (CA). **JOLICOEUR, Eric** [CA/CA]; 2100 Cunard Street, Laval, Québec H7S 2G5 (CA). **GILLARD, James** [CA/CA]; 2100 Cunard Street, Laval, Québec H7S 2G5 (CA). **POUPART, Marc-André** [CA/CA]; 2100 Cunard Street, Laval, Québec H7S 2G5 (CA). **RANCOURT, Jean**


Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/010140 A2

(54) Title: **VIRAL POLYMERASE INHIBITORS**



definition as R⁴ defined above; M is N or CR⁷, wherein R⁷ has the same definition as R⁴ defined above; Y¹ is O or S; Z is N(R^{6a})R⁶ or OR⁶, wherein R^{6a} is H or alkyl or NR⁶¹R⁶² wherein R⁶¹ and R⁶² are defined herein; a salt or a derivative thereof, as an inhibitor of HCV NS^{5B} polymerase.

(57) Abstract: An isomer, enantiomer, diastereoisomer, or tautomer of a compound, represented by formula (I): wherein: A is O, S, NR¹, or CR₁, wherein R₁ is defined herein; represents either a single or a double bond; R₂ is selected from: H, halogen, R₂₁, OR₂₁, SR₂₁, COOR₂₁, SO₂N(R²²)₂, N(R²²)₂, CON(R²²)₂, NR²²C(O)R²² or NR²²C(O)NR²² wherein R²¹ and each R²² is defined herein; B is NR³ or CR³, with the proviso that one of A or B is either CR¹ or CR³, wherein R³ is defined herein; K is N or CR⁴, wherein R⁴ is defined herein; L is N or CR⁵, wherein R⁵ has the same

VIRAL POLYMERASE INHIBITORS

TECHNICAL FIELD OF THE INVENTION

The invention relates to inhibitors of RNA dependent RNA polymerases,
5 particularly those viral polymerases within the Flaviviridae family, more particularly to HCV polymerase.

BACKGROUND OF THE INVENTION

About 30,000 new cases of hepatitis C virus (HCV) infection are estimated to
10 occur in the United States each year (Kolykhalov, A.A.; Mihalik, K.; Feinstone, S.M.; Rice, C.M.; 2000; *J. Virol.* **74**: 2046-2051). HCV is not easily cleared by the hosts' immunological defences; as many as 85% of the people infected with HCV become chronically infected. Many of these persistent infections result in chronic liver disease, including cirrhosis and hepatocellular carcinoma (Hoofnagle, J.H.;
15 1997; *Hepatology* **26**: 15S-20S). There are an estimated 170 million HCV carriers world-wide, and HCV-associated end-stage liver disease is now the leading cause of liver transplantation. In the United States alone, hepatitis C is responsible for 8,000 to 10,000 deaths annually. Without effective intervention, the number is expected to triple in the next 10 to 20 years. There is no vaccine to
20 prevent HCV infection. Prolonged treatment of chronically infected patients with interferon or interferon and ribavirin is the only currently approved therapy, but it achieves a sustained response in fewer than 50% of cases (Lindsay, K.L.; 1997; *Hepatology* **26**: 71S-77S, and Reichard, O.; Schvarcz, R.; Weiland, O.; 1997 *Hepatology* **26**: 108S-111S).

25

HCV belongs to the family *Flaviviridae*, genus *hepacivirus*, which comprises three genera of small enveloped positive-strand RNA viruses (Rice, C.M.; 1996; "Flaviviridae: the viruses and their replication"; pp. 931-960 in *Fields Virology*,
Fields, B.N.; Knipe, D.M.; Howley, P.M. (eds.); Lippincott-Raven Publishers,
30 Philadelphia Pa.). The 9.6 kb genome of HCV consists of a long open reading frame (ORF) flanked by 5' and 3' non-translated regions (NTR's). The HCV 5' NTR is 341 nucleotides in length and functions as an internal ribosome entry site for cap-independent translation initiation (Lemon, S.H.; Honda, M.; 1997; *Semin. Virol.* **8**: 274-288). The HCV polyprotein is cleaved co- and post-translationally
35 into at least 10 individual polypeptides (Reed, K.E.; Rice, C.M.; 1999; *Curr. Top.*

Microbiol. Immunol. **242**: 55-84). The structural proteins result from signal peptidases in the N-terminal portion of the polyprotein. Two viral proteases mediate downstream cleavages to produce non-structural (NS) proteins that function as components of the HCV RNA replicase. The NS2-3 protease spans the C-terminal half of the NS2 and the N-terminal one-third of NS3 and catalyses *cis* cleavage of the NS2/3 site. The same portion of NS3 also encodes the catalytic domain of the NS3-4A serine protease that cleaves at four downstream sites. The C-terminal two-thirds of NS3 is highly conserved amongst HCV isolates, with RNA-binding, RNA-stimulated NTPase, and RNA unwinding activities. Although NS4B and the NS5A phosphoprotein are also likely components of the replicase, their specific roles are unknown. The C-terminal polyprotein cleavage product, NS5B, is the elongation subunit of the HCV replicase possessing RNA-dependent RNA polymerase (RdRp) activity (Behrens, S.E.; Tomei, L.; DeFrancesco, R.; 1996; *EMBO J.* **15**: 12-22; and Lohmann, V.; Körner, F.; Herian, U.; Bartenschlager, R.; 1997; *J. Virol.* **71**: 8416-8428). It has been recently demonstrated that mutations destroying NS5B activity abolish infectivity of RNA in a chimp model (Kolykhalov, A.A.; Mihalik, K.; Feinstone, S.M.; Rice, C.M.; 2000; *J. Virol.* **74**: 2046-2051).

The development of new and specific anti-HCV treatments is a high priority, and virus-specific functions essential for replication are the most attractive targets for drug development. The absence of RNA dependent RNA polymerases in mammals, and the fact that this enzyme appears to be essential to viral replication, would suggest that the NS5B polymerase is an ideal target for anti-HCV therapeutics.

WO 00/06529 reports inhibitors of NS5B which are α , γ -diketoacids.

WO 00/13708, WO 00/10573, WO 00/18231, and WO 01/47883 report inhibitors of NS5B proposed for treatment of HCV.


SUMMARY OF THE INVENTION

It is therefore an object of the invention to provide a novel series of compounds having improved inhibitory activity against HCV polymerase.

In a first aspect of the invention, there is provided an isomer, enantiomer,

3

diastereoisomer, or tautomer of a compound, represented by formula I:



wherein:

A is O, S, NR¹, or CR¹, wherein R¹ is selected from the group consisting of: H,

5 (C₁₋₆)alkyl optionally substituted with:

-halogen, OR¹¹, SR¹¹ or N(R¹²)₂, wherein R¹¹ and each R¹² is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-aryl or (C₁₋₆)alkyl-Het, said aryl or Het optionally substituted with R¹⁰; or

10 both R¹² are covalently bonded together and to the nitrogen to which they are both attached to form a 5, 6 or 7-membered saturated heterocycle;

----- represents either a single or a double bond;

15 R² is selected from: H, halogen, R²¹, OR²¹, SR²¹, COOR²¹, SO₂N(R²²)₂, N(R²²)₂, CON(R²²)₂, NR²²C(O)R²² or NR²²C(O)NR²² wherein R²¹ and each R²² is independently H, (C₁₋₆)alkyl, haloalkyl, (C₂₋₆)alkenyl, (C₃₋₇)cycloalkyl, (C₂₋₆)alkynyl, (C₅₋₇)cycloalkenyl, 6 or 10-membered aryl or Het, said R²¹ and R²² being optionally substituted with R²⁰, or both R²² are bonded together to form a 5, 6 or 7-

20 membered saturated heterocycle with the nitrogen to which they are attached; wherein R¹⁰ and R²⁰ is each:

- 1 to 4 substituents selected from: halogen, OPO₃H, NO₂, cyano, azido, C(=NH)NH₂, C(=NH)NH(C₁₋₆)alkyl or C(=NH)NHCO(C₁₋₆)alkyl; or

- 1 to 4 substituents selected from:

25 a) (C₁₋₆) alkyl or haloalkyl, (C₃₋₇)cycloalkyl, C₃₋₇ spirocycloalkyl optionally containing 1 or 2 heteroatom, (C₂₋₆)alkenyl, (C₃₋₆)cycloalkenyl, (C₂₋₈)alkynyl, (C₁₋₆) alkyl-(C₃₋₇)cycloalkyl, all of which optionally substituted with R¹⁵⁰;

b) OR¹⁰⁴ wherein R¹⁰⁴ is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, Het, (C₁₋₆)alkyl)aryl or (C₁₋₆)alkyl)Het, said alkyl, cycloalkyl, aryl, Het, (C₁₋₆)alkyl)aryl or (C₁₋₆)alkyl)Het being optionally substituted with

30 R¹⁵⁰;

- c) OCOR^{105} wherein R^{105} is (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het**, said alkyl, cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het** being optionally substituted with R^{150} ;
- d) SR^{108} , $\text{SO}_2\text{N}(\text{R}^{108})_2$ or $\text{SO}_2\text{N}(\text{R}^{108})\text{C}(\text{O})\text{R}^{108}$ wherein each R^{108} is independently H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl or (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het** or both R^{108} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het** or heterocycle being optionally substituted with R^{150} ;
- e) $\text{NR}^{111}\text{R}^{112}$ wherein R^{111} is H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl or (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het**, and R^{112} is H, CN, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl or (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl, (C_{1-6}) alkyl)**Het**, COOR^{115} or $\text{SO}_2\text{R}^{115}$ wherein R^{115} is (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, or (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het**, or both R^{111} and R^{112} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het**, or heterocycle being optionally substituted with R^{150} ;
- f) $\text{NR}^{116}\text{COR}^{117}$ wherein R^{116} and R^{117} is each H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het**, said (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het** being optionally substituted with R^{150} ;
- g) $\text{NR}^{118}\text{CONR}^{119}\text{R}^{120}$, wherein R^{118} , R^{119} and R^{120} is each H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het**, or R^{118} is covalently bonded to R^{119} and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle; or R^{119} and R^{120} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het** or heterocycle being optionally substituted with R^{150} ;
- h) $\text{NR}^{121}\text{COCOR}^{122}$ wherein R^{121} and R^{122} is each H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, a 6- or 10-membered aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het** or heterocycle being optionally substituted with R^{150} ;

5

- alkyl)aryl or (C₁₋₆alkyl)Het, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, Het, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)Het being optionally substituted with R¹⁵⁰; or R¹²² is OR¹²³ or N(R¹²⁴)₂ wherein R¹²³ and each R¹²⁴ is independently H, (C₁₋₆alkyl), (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, Het, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)Het, or R¹²⁴ is OH or O(C₁₋₆alkyl) or both R¹²⁴ are covalently bonded together to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, Het, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)Het and heterocycle being optionally substituted with R¹⁵⁰;
- i) COR¹²⁷ wherein R¹²⁷ is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, Het, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)Het, said alkyl, cycloalkyl, aryl, Het, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)Het being optionally substituted with R¹⁵⁰;
- j) COOR¹²⁸ wherein R¹²⁸ is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, Het, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)Het, said (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, Het, (C₁₋₆alkyl)aryl and (C₁₋₆alkyl)Het being optionally substituted with R¹⁵⁰;
- k) CONR¹²⁹R¹³⁰ wherein R¹²⁹ and R¹³⁰ are independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, Het, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)Het, or both R¹²⁹ and R¹³⁰ are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, Het, (C₁₋₆alkyl)aryl, (C₁₋₆alkyl)Het and heterocycle being optionally substituted with R¹⁵⁰;
- l) aryl, Het, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)Het, all of which being optionally substituted with R¹⁵⁰; wherein R¹⁵⁰ is defined as:
- 1 to 3 substituents selected from: halogen, OPO₃H, NO₂, cyano, azido, C(=NH)NH₂, C(=NH)NH(C₁₋₆)alkyl or C(=NH)NHCO(C₁₋₆)alkyl; or
 - 1 to 3 substituents selected from:
- a) (C₁₋₆) alkyl or haloalkyl, (C₃₋₇)cycloalkyl, C₃₋₇ spirocycloalkyl optionally containing 1 or 2 heteroatom, (C₂₋₆)alkenyl, (C₂₋₈)alkynyl, (C₁₋₆) alkyl-(C₃₋₇)cycloalkyl, all of which optionally substituted with R¹⁶⁰;
- b) OR¹⁰⁴ wherein R¹⁰⁴ is H, (C₁₋₆alkyl), (C₃₋₇)cycloalkyl, or (C₁₋

₆alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said alkyl, cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being optionally substituted with R¹⁶⁰;

5

c) OCOR¹⁰⁵ wherein R¹⁰⁵ is (C₁₋₆alkyl), (C₃₋₇)cycloalkyl, (C₁₋₆alkyl)-(C₃₋₇)cycloalkyl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said alkyl, cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being optionally substituted with R¹⁶⁰;

10

d) SR¹⁰⁸, SO₂N(R¹⁰⁸)₂ or SO₂N(R¹⁰⁸)C(O)R¹⁰⁸ wherein each R¹⁰⁸ is independently H, (C₁₋₆alkyl), (C₃₋₇)cycloalkyl or (C₁₋₆alkyl)-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** or both R¹⁰⁸ are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** or heterocycle being optionally substituted with R¹⁶⁰;

15

e) NR¹¹¹R¹¹² wherein R¹¹¹ is H, (C₁₋₆alkyl), (C₃₋₇)cycloalkyl or (C₁₋₆alkyl)-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, and R¹¹² is H, CN, (C₁₋₆alkyl), (C₃₋₇)cycloalkyl or (C₁₋₆alkyl)-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl, (C₁₋₆alkyl)**Het**, COOR¹¹⁵ or SO₂R¹¹⁵ wherein R¹¹⁵ is (C₁₋₆alkyl), (C₃₋₇)cycloalkyl, or (C₁₋₆alkyl)-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, or both R¹¹¹ and R¹¹² are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, or heterocycle being optionally substituted with R¹⁶⁰;

20

25

f) NR¹¹⁶COR¹¹⁷ wherein R¹¹⁶ and R¹¹⁷ is each H, (C₁₋₆alkyl), (C₃₋₇)cycloalkyl, (C₁₋₆alkyl)-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said (C₁₋₆alkyl), (C₃₋₇)cycloalkyl, (C₁₋₆alkyl)-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being optionally substituted with R¹⁶⁰;

30

g) NR¹¹⁸CONR¹¹⁹R¹²⁰, wherein R¹¹⁸, R¹¹⁹ and R¹²⁰ is each H, (C₁₋₆alkyl), (C₃₋₇)cycloalkyl, (C₁₋₆alkyl)-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, or R¹¹⁸ is covalently bonded to R¹¹⁹ and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, or R¹¹⁹ and R¹²⁰ are covalently

7

- bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** or heterocycle being optionally substituted with R¹⁶⁰;
- 5 **h) NR¹²¹COCOR¹²²** wherein R¹²¹ and R¹²² is each H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, a 6- or 10-membered aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being optionally substituted with R¹⁶⁰, or R¹²² is OR¹²³ or N(R¹²⁴)₂ wherein
- 10 R¹²³ and each R¹²⁴ is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, or R¹²⁴ is OH or O(C₁₋₆alkyl) or both R¹²⁴ are covalently bonded together to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** and heterocycle being optionally
- 15 substituted with R¹⁶⁰;
- i) COR¹²⁷** wherein R¹²⁷ is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said alkyl, cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being
- 20 optionally substituted with R¹⁶⁰;
- j) tetrazole, COOR¹²⁸** wherein R¹²⁸ is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl and (C₁₋₆alkyl)**Het** being
- 25 optionally substituted with R¹⁶⁰; and
- k) CONR¹²⁹R¹³⁰** wherein R¹²⁹ and R¹³⁰ are independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, or both R¹²⁹ and R¹³⁰ are covalently bonded together and to the nitrogen to which they are attached to
- 30 form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl, (C₁₋₆alkyl)**Het** and heterocycle being optionally substituted with R¹⁶⁰;
- wherein R¹⁶⁰ is defined as 1 or 2 substituents selected from: tetrazole, halogen, CN, C₁₋₆alkyl, haloalkyl, COOR¹⁶¹,

8

5 SO_3H , SR^{161} , $\text{SO}_2\text{R}^{161}$, OR^{161} , $\text{N}(\text{R}^{162})_2$, $\text{SO}_2\text{N}(\text{R}^{162})_2$,
 $\text{NR}^{162}\text{COR}^{162}$ or $\text{CON}(\text{R}^{162})_2$, wherein R^{161} and each R^{162} is
independently H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl or (C_{1-6}) alkyl-
 (C_{3-7}) cycloalkyl; or both R^{162} are covalently bonded together
and to the nitrogen to which they are attached to form a 5,
6 or 7-membered saturated heterocycle;

B is NR^3 or CR^3 , with the proviso that one of **A** or **B** is either CR^1 or CR^3 ,
wherein R^3 is selected from (C_{1-6}) alkyl; haloalkyl, (C_{3-7}) cycloalkyl, (C_{5-7}) cycloalkenyl,
10 (C_{6-10}) bicycloalkyl, (C_{6-10}) bicycloalkenyl, 6- or 10-membered aryl,
Het, (C_{1-6}) alkyl-aryl or (C_{1-6}) alkyl-**Het**,
said alkyl, cycloalkyl, bicycloalkyl, aryl, **Het**, alkyl-aryl and alkyl-**Het** being
optionally substituted with from 1 to 4 substituents selected from: halogen,
or

- 15 a) (C_{1-6}) alkyl optionally substituted with:
- OR^{31} or SR^{31} wherein R^{31} is H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl,
 (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl-aryl or (C_{1-6}) alkyl-**Het**; or
 - $\text{N}(\text{R}^{32})_2$ wherein each R^{32} is independently H, (C_{1-6}) alkyl,
20 (C_{3-7}) cycloalkyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl-aryl or (C_{1-6}) alkyl-**Het**; or both R^{32} are covalently
bonded together and to the nitrogen to which they are
attached to form a 5, 6 or 7-membered saturated
heterocycle;
- 25 b) OR^{33} wherein R^{33} is H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl or
 (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl-aryl or (C_{1-6}) alkyl-**Het**;
- c) SR^{34} wherein R^{34} is H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, or
 (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl-aryl or (C_{1-6}) alkyl-**Het**; and
- 30 d) $\text{N}(\text{R}^{35})_2$ wherein each R^{35} is independently H, (C_{1-6}) alkyl,
 (C_{3-7}) cycloalkyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl-aryl
or (C_{1-6}) alkyl-**Het**; or both R^{35} are covalently bonded together and
to the nitrogen to which they are attached to form a 5, 6 or 7-

membered saturated heterocycle;

K is N or CR⁴, wherein R⁴ is H, halogen, (C₁₋₆)alkyl, haloalkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl; or R⁴ is OR⁴¹ or SR⁴¹, COR⁴¹ or NR⁴¹COR⁴¹ wherein
 5 each R⁴¹ is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, or R⁴ is NR⁴²R⁴³ wherein R⁴² and R⁴³ are each independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, or both R⁴² and R⁴³ are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle;

10

L is N or CR⁵, wherein R⁵ has the same definition as R⁴ defined above;


M is N or CR⁷, wherein R⁷ has the same definition as R⁴ defined above;

15 Y¹ is O or S;

Z is OR⁶, wherein R⁶ is H, (C₁₋₆)alkyl being optionally substituted with: halo, hydroxy, carboxy, amino, C₁₋₆ alkoxy, C₁₋₆alkoxycarbonyl, and C₁₋₆ alkylamino; or R⁶ is C₁₋₆ alkylaryl optionally substituted with: halogen, cyano, nitro, C₁₋₆ alkyl, C₁₋₆haloalkyl, C₁₋₆alkanoyl, -(CH₂)₁₋₆-COOR⁷, -(CH₂)₁₋₆-CONR⁷R⁸, -(CH₂)₁₋₆-NR⁷R⁸, -
 20 (CH₂)₁₋₆-NR⁷COR⁸, -(CH₂)₁₋₆-NH₂SO₂R⁷, -(CH₂)₁₋₆-OR⁷, -(CH₂)₁₋₆-SR⁷, -(CH₂)₁₋₆-SO₂R⁷, and -(CH₂)₁₋₆-SO₂NR⁷R⁸, wherein each R⁷ and each R⁸ is H or C₁₋₆ alkyl,

or Z is NR⁹R¹⁰ wherein each of R⁹ and R¹⁰ is selected from: H, C₁₋₆alkoxy, or C₁₋₆alkyl optionally substituted with halo, hydroxy, carboxy, amino, C₁₋₆ alkoxy, C₁₋₆alkoxycarbonyl, and C₁₋₆ alkylamino;
 25

or a salt thereof;

30 with the proviso that when A is CR¹, R¹ is Me, R² is pyridine or , B is NR³, R³ is Me, K, L, M is CH, Y¹ is O, and Z is OR⁶, then R⁶ is not H;

and with the proviso that when A is NR¹, R¹ is H, R² is phenyl, B is CR³, R³ is

10

phenyl, **K**, **L**, **M** is CH, **Y**¹ is O, and **Z** is OR⁶, then **R**⁶ is not H;


and with the proviso that when **A** is S, **R**² is bromine, **B** is CR³, **R**³ is Me, **K** is CH, **L** is CH, **M** is CR⁷, **R**⁷ is H or Me, **Y**¹ is O, and **Z** is OR⁶, then **R**⁶ is not H;

5

and with the proviso that when **A** is O, **R**² is H, **B** is CR³, **R**³ is phenyl, **K**, **L**, **M** is CH, **Y**¹ is O, and **Z** is OR⁶, then **R**⁶ is not H;

and with the proviso that when **A** is CR¹, **R**¹ is Me, **R**² is pyridine, **B** is NR³, **R**³ is Me, **K**, **L**, **M** is CH, **Y**¹ is O, and **Z** is OR⁶, then **R**⁶ is not Me;

10

and with the further proviso that when **A** is CR¹, **R**¹ is Me, **R**² is , **B** is NR³, **R**³ is Me, **K**, **L**, **M** is CH, **Y**¹ is O, and **Z** is OR⁶, then **R**⁶ is not Et;


and with the further proviso that when **A** is CR¹, **R**¹ is CH, **R**² is Me, **B** is NR³, **R**³ is Me, **K**, **L**, **M** is CH, **Y**¹ is O, and **Z** is OR⁶, then **R**⁶ is not Et;

15

and with the further proviso that when **A** is CR¹, **R**¹ is Et, **R**² is Me, **B** is NR³, **R**³ is Me, **K**, **L**, **M** is CH, **Y**¹ is O, and **Z** is OR⁶, then **R**⁶ is not CH₂CH₂N(Me)₂;

20

and with the further proviso that when **A** is CH, **R**² is Me, **B** is NR³, **R**³ is




, **K** is N, **L** is CR⁵, **R**⁵ is Me, **M** is CR⁷, **R**⁷ is OH, **Y**¹ is O, and **Z** is OR⁶ then **R**⁶ is not Et;

and with the further proviso that when **A** is NR¹, **R**¹ is Me, **R**² is Br, **B** is CR³, **R**³ is

25


11




, K is N, L is CR⁵, R⁵ is Me, M is CR⁷, R⁷ is Br, Y¹ is O, and Z is OR⁶, then R⁶ is not Me;

and with the further proviso that when A is NR¹, R¹ is H, R² is Cl, B is CR³, R³ is Et, K is CH, L is CH, M is CH, Y¹ is O, Z is OR⁶, then R⁶ is not Me;

and with the further proviso that when A is NR¹, R¹ is H, R² is phenyl, B is CR³, R³ is phenyl, K is CH, L is CH, M is CR⁷, R⁷ is Me, Y¹ is O, Z is OR⁶, then R⁶ is not Et;



10 and with the further proviso that when A is NR¹, R¹ is H, R² is OMe, B is CR³,




R³ is OMe, K is CH, L is N, M is CH, Y¹ is O, and Z is OR⁶, then R⁶ is not Et;

with the further proviso that when A is S, R² is Br, B is CR³, R³ is Me, K is CH, L is CH, M is CH, Y¹ is O, and Z is OR⁶, then R⁶ is not Me;


15

and with the further proviso that, when A is NR¹, R¹ is H, R² is:



, B is NR³, R³ is cyclohexyl, K, L, M is CH, Y¹ is O, Z is OR⁶, then R⁶ is not H.

Alternatively, in a first aspect of the invention, there is provided a compound represented by Formula Ia:



wherein:

A is O, S, NR¹, or CR¹;

5 B is NR³ or CR³;

R¹ is selected from the group consisting of: H, (C₁₋₆)alkyl, benzyl, (C₁₋₆ alkyl)-(C₆₋₁₀aryl), (C₁₋₆ alkyl)-5- or 6-membered heterocycle having 1 to 4 heteroatoms selected from O, N, and S, and 5- or 6-membered heterocycle having 1 to 4
10 heteroatoms selected from O, N and S,

wherein said benzyl and said heteroatom are optionally substituted with from 1 to 4 substituents selected from the group consisting of: COOH, COO(C₁₋₆ alkyl), halogen, and (C₁₋₆ alkyl);

15 R² is selected from the group consisting of: H, halogen, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, phenyl, 5- or 6-membered heterocycle having 1 to 4 heteroatoms selected from O, N, and S, pyridine-N-oxide, and 9- or 10-membered heterobicyclic having 1 to 4 heteroatoms selected from O, N and S,

20 said phenyl, heterocycle and heterobicyclic being optionally substituted with from 1 to 4 substituents selected from the group consisting of: halogen, C(halogen)₃, (C₁₋₆)alkyl, OH, O(C₁₋₆ alkyl), NH₂, and N(C₁₋₆ alkyl)₂;

R³ is selected from the group consisting of: 5-, 6- or 7-membered heterocycle having 1 to 4 heteroatoms selected from O, N, and S, norbornane, (C₃₋₇)cycloalkyl and (C₃₋₇)cycloalkyl-(C₁₋₆ alkyl);
25

M is N, CR⁴, or COR⁵, wherein R⁴ is selected from the group consisting of: H, halogen, and (C₁₋₆ alkyl); and R⁵ is selected from the group consisting of: H and (C₁₋₆ alkyl);

30

K and L is N or CH;

----- represents either a single or a double bond;

Y is O;

5

Z is OR^6 or NR^6R^{6a} , wherein R^6 is selected from the group consisting of: H, (C_{1-6}) alkyl, wherein said alkyl is optionally substituted with from 1 to 4 substituents selected from: OH, COOH, $COO(C_{1-6})$ alkyl, (C_{1-6}) alkyl, said alkyl being optionally substituted with from 1 to 4 substituents selected from: COOH, $NHCO(C_{1-6})$ alkyl, 10 NH_2 , $NH(C_{1-6})$ alkyl, and $N(C_{1-6})_2$ alkyl;

or a salt thereof.

15 In a third aspect of the invention, there is provided a compound of the formula I, or a pharmaceutically acceptable salt thereof, as an inhibitor of RNA dependent RNA polymerase activity of the enzyme NS5B, encoded by HCV.

20 In a fourth aspect of the invention, there is provided a compound of the formula I, or a pharmaceutically acceptable salt thereof, as an inhibitor of HCV replication.

25 In a fifth aspect of the invention, there is provided a method of treating or preventing HCV infection in a mammal, comprising administering to the mammal an effective amount of a compound of formula I, or a pharmaceutically acceptable salt thereof.

30 In a sixth aspect of the invention, there is provided a pharmaceutical composition for the treatment or prevention of HCV infection, comprising an effective amount of a compound of formula I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

35 According to a specific embodiment, the pharmaceutical compositions of this invention comprise an additional immunomodulatory agent. Examples of additional immunomodulatory agents include but are not limited to, α -, β -, δ -, γ -, and ω -interferons.

14

According to an alternate embodiment, the pharmaceutical compositions of this invention may additionally comprise an antiviral agent. Examples of antiviral agents include, ribavirin and amantadine.

- 5 According to another alternate embodiment, the pharmaceutical compositions of this invention may additionally comprise other inhibitors of HCV polymerase.


According to yet another alternate embodiment, the pharmaceutical compositions of this invention may additionally comprise an inhibitor of other targets in the HCV
10 life cycle, such as helicase, polymerase, metalloprotease or IRES.

In a seventh aspect of the invention, there is provided a use of a compound of formula I, for the manufacture of a medicament for the treatment of HCV infection.

- 15 In an eighth aspect of the invention, there is provided a use of a compound of formula I, as an HCV polymerase inhibitor.

In a ninth aspect of the invention, there is provided a method of treating or preventing HCV infection in a mammal, comprising administering to the mammal
20 an effective amount of a compound of formula I, or a pharmaceutically acceptable salt thereof in combination with another anti-HCV agent.


In a tenth aspect of the invention, there is provided an intermediate of formula (1a) or (1b):



wherein **A**, **B**, **K**, **L**, and **M** are as described herein and **PG** is H or a carboxy protecting group.

- 25 In a eleventh aspect of the invention, there is provided the use of the intermediates of formula (1a) for producing compounds of formula (iii),


15



(iii)

wherein **A**, **R²**, **B**, **K**, **L**, **M**, and **PG** are as described herein,
comprising:


- a) coupling, in the presence of a metal catalyst (such as, for example, Pd, Ni,
Ru, Cu), a base and an additive (such as a phosphine ligand, Cu salt, Li salt,
ammonium salt, CsF) in an appropriate solvent, intermediate (1a):



1a


- with **R²-X**, wherein **R¹**, **R³**, **K**, **L**, **M** and **PG** are as described herein and **X** is (bt
not limited to): Sn(C₁₋₆alkyl)₃, Sn(aryl)₃, metal halide, B(OH)₂, and B(O(C₁₋₆alkyl)₂)
to produce compounds of formula (iii).

In an alternative to the eleventh aspect of the invention, there is provided the use
of intermediate (1b) for producing compounds of formula (iii),



(iii)

- wherein **A**, **R²**, **B**, **K**, **L**, **M**, and **PG** are as described herein,
comprising:
b) coupling, in the presence of a metal catalyst (such as, for example, Pd, Ni,
Ru, Cu), a base and an additive (such as a phosphine ligand, Cu salt, Li salt,
ammonium salt, CsF) in an appropriate solvent, intermediate (1b)



1b

with **R²-X'**, wherein **X'** is halide, OSO₂(C₁₋₆alkyl), OSO₂Ar, OSO₂CF₃ and the like,

16

and **M** is a metal such as Li, Sn(C₁₋₆alkyl)₃, Sn(aryl)₃, B(OH)₂, B(OC₁₋₆alkyl)₂, metal halide, to produce compounds of formula (iii).

DETAILED DESCRIPTION OF THE INVENTION5 **Definitions**

The following definitions apply unless otherwise noted:

As used herein, the terms "(C₁₋₃) alkyl", "(C₁₋₄) alkyl" or "(C₁₋₆) alkyl", either alone or in combination with another radical, are intended to mean acyclic straight or
10 branched chain alkyl radicals containing up to three, four and six carbon atoms respectively. Examples of such radicals include methyl, ethyl, propyl, butyl, hexyl, 1-methylethyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl.

As used herein, the term "(C₂₋₆) alkenyl", either alone or in combination with
15 another radical, is intended to mean an unsaturated, acyclic straight chain radical containing two to six carbon atoms.

As used herein, the term "(C₂₋₆) alkynyl" either alone or in combination with another group, is intended to mean an unsaturated, acyclic straight chain sp hybridized
20 radical containing 2 to six carbon atoms.

As used herein, the term "(C₃₋₇) cycloalkyl", either alone or in combination with another radical, means a cycloalkyl radical containing from three to seven carbon atoms and includes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and
25 cycloheptyl.

As used herein, the term "(C₅₋₇)cycloalkenyl", either alone or in combination with another radical, means an unsaturated cyclic radical containing five to seven carbon atoms.

30


As used herein, the term "carboxy protecting group" defines protecting groups that can be used during coupling and are listed in Greene, "Protective Groups in Organic Chemistry", John Wiley & Sons, New York (1981) and "The Peptides: Analysis, Synthesis, Biology", Vol. 3, Academic Press, New York (1981), the
35 disclosures of which are hereby incorporated by reference.

The α -carboxyl group of the C-terminal residue is usually protected as an ester (CPG) that can be cleaved to give the carboxylic acid. Protecting groups that can be used include: 1) alkyl esters such as methyl, trimethylsilylethyl and *t*-butyl, 2)
5 aralkyl esters such as benzyl and substituted benzyl, or 3) esters that can be cleaved by mild base treatment or mild reductive means such as trichloroethyl and phenacyl esters.

As used herein, the term "aryl", or "6- or 10-membered aryl" either alone or in
10 combination with another radical means aromatic radical containing six or ten carbon atoms, for example phenyl or naphthyl.


As used herein the term heteroatom means O, S or N.

15 As used herein, the term "heterocycle", either alone or in combination with another radical, means a monovalent radical derived by removal of a hydrogen from a five-, six-, or seven-membered saturated or unsaturated (including aromatic) heterocycle containing from one to four heteroatoms selected from nitrogen, oxygen and sulfur. Furthermore, "heterobicyclic" as used herein, means
20 a heterocycle as defined above fused to one or more other cycle, be it a heterocycle or any other cycle. Examples of such heterocycles include, but are not limited to, pyrrolidine, tetrahydrofuran, thiazolidine, pyrrole, thiophene, coumarin, hydantoin, diazepine, 1H-imidazole, isoxazole, thiazole, tetrazole, piperidine, 1,4-dioxane, 4-morpholine, pyridine, pyridine-N-oxide, pyrimidine,
25 thiazolo[4,5-b]-pyridine, quinoline, or indole, or the following heterocycles:



As used herein, the term "9- or 10-membered heterobicyclic" or "heterobicyclic" either alone or in combination with another radical, means a heterocycle as
30 defined above fused to one or more other cycle, be it a heterocycle or any other cycle. Examples of such heterobicyclics include, but are not limited to, thiazolo[4,5-b]-pyridine, quinoline, or indole, or the following:

18



As used herein, the term "Het" defines a 5- or 6-membered heterocycle having 1
 5 to 4 heteroatoms selected from O, N, and S, or a 9- or 10-membered
 heterobicyclic having 1 to 5 heteroatoms wherever possible, selected from O, N
 and S.

As used herein, the term "halo" means a halogen atom and includes fluorine,
 10 chlorine, bromine and iodine.

As used herein, the term "haloalkyl" is intended to mean an alkyl that is described
 above in which each hydrogen atom may be successively replaced by a halogen
 atom, for example CH_2Br or CF_3 .

15

As used herein, the term "metal halide" is intended to mean any metal that is
 bonded to a halogen atom for use in a metal-catalyzed cross-coupling reaction.
 Examples of such metal halides include, but are not limited to, $-\text{MgCl}$, $-\text{CuCl}$, or $-\text{ZnCl}$
 and the like.

20

As used herein, the term "OH" refers to a hydroxyl group. It is well known to one
 skilled in the art that hydroxyl groups may be substituted by functional group
 equivalents. Examples of such functional group equivalents that are
 contemplated by this invention include, but are not limited to, ethers, sulfhydryls,
 25 and primary, secondary or tertiary amines.

As used herein, the term "SH" refers to a sulfhydryl group. It is intended within the
 scope of the present invention that, whenever a "SH" or "SR" group is present, it
 can also be substituted by any other appropriate oxidation state such as SOR ,
 30 SO_2R , or SO_3R .

It is intended that the term "substituted" when applied in conjunction with a radical having more than one moiety such as C₁₋₆alkyl-aryl, or C₁₋₆alkyl-Het, such substitution applies to both moieties i.e. both the alkyl and aryl or Het moieties
5 can be substituted with the defined substituents.

As used herein, the term "COOH" refers to a carboxylic acid group. It is well known to one skilled in the art that carboxylic acid groups may be substituted by functional group equivalents. Examples of such functional group equivalents that
10 are contemplated by this invention include, but are not limited to, esters, amides, boronic acids or tetrazole.

As used herein, the term "functional group equivalent" is intended to mean an element or a substituted derivative thereof, that is replaceable by another element
15 that has similar electronic, hybridization or bonding properties.

As used herein, the term "metal catalyst" is intended to mean a metal such as palladium (0) or palladium (2) that is bonded to a leaving group for use in a cross-coupling reaction. Examples of such palladium catalysts include, but are not
20 limited to, Pd(Ph₃)₄, Pd/C, Pd(OAc)₂, PdCl₂, and the like. Alternative metals that can catalyze cross-coupling reactions include, but are not limited to: Ni(acac)₂, Ni(OAc)₂, or NiCl₂.

As used herein, the term "derivative" is intended to mean "detectable label",
25 "affinity tag" or "photoreactive group". The term "detectable label" refers to any group that may be linked to the polymerase or to a compound of the present invention such that when the compound is associated with the polymerase target, such label allows recognition either directly or indirectly of the compound such that it can be detected, measured and quantified. Examples of such "labels" are
30 intended to include, but are not limited to, fluorescent labels, chemiluminescent labels, colorimetric labels, enzymatic markers, radioactive isotopes and affinity tags such as biotin. Such labels are attached to the compound or to the polymerase by well known methods.

The term "affinity tag" means a ligand (that is linked to the polymerase or to a
35 compound of the present invention) whose strong affinity for a receptor can be

used to extract from a solution the entity to which the ligand is attached. Examples of such ligands include biotin or a derivative thereof, a histidine polypeptide, a polyarginine, an amylose sugar moiety or a defined epitope recognizable by a specific antibody. Such affinity tags are attached to the compound or to the polymerase by well-known methods.


The term "photoreactive group" means a group that is transformed, upon activation by light, from an inert group to a reactive species, such as a free radical. Examples of such groups include, but are not limited to, benzophenones, azides, and the like.

As used herein, the term "pharmaceutically acceptable salt" includes those derived from pharmaceutically acceptable bases and is non-toxic. Examples of suitable bases include choline, ethanolamine and ethylenediamine. Na^+ , K^+ , and Ca^{++} salts are also contemplated to be within the scope of the invention (also see Pharmaceutical salts, Birge, S.M. et al., J. Pharm. Sci., (1977), 66, 1-19, incorporated herein by reference).

Preferred embodiments

A:

Preferably, compounds of the present invention have the following formula (II):




wherein, preferably, A is O, S, or NR^1 .

Preferably, A is NR^1 .

Preferably, compounds of the present invention have the following formula (III):


21




wherein, preferably, **B** is NR^3 .

With respect to compounds of formula (II) and (III), preferably, **M**, **K** and **L** is CH
 5 or N. More preferably, **M**, **K** and **L** is CH.


More preferably, compounds of the present invention have the following formulae:




IIa




IIIa



IIb



IIc



or

IIld

15 R^1 :

Preferably R^1 is selected from the group consisting of: H or (C_{1-6}) alkyl. More preferably, R^1 is H, CH_3 , isopropyl, or isobutyl. Even more preferably, R^1 is H or CH_3 . Most preferably, R^1 is CH_3 .

20 R^2 :

Preferably, R^2 is selected from: H, halogen, (C_{2-6}) alkenyl, (C_{5-7}) cycloalkenyl, 6 or 10-membered aryl or **Het**; wherein (C_{2-6}) alkenyl, (C_{5-7}) cycloalkenyl, aryl or **Het** is optionally substituted with R^{20} , wherein R^{20} is defined as:

- 1 to 4 substituents selected from: halogen, NO_2 , cyano, azido,
 5 $C(=NH)NH_2$, $C(=NH)NH(C_{1-6})$ alkyl or $C(=NH)NHCO(C_{1-6})$ alkyl; or
- 1 to 4 substituents selected from:
 - a) (C_{1-6}) alkyl or haloalkyl, (C_{3-7}) cycloalkyl, (C_{2-6}) alkenyl, (C_{2-8}) alkynyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, all of which optionally substituted with R^{150} ;
 - b) OR^{104} wherein R^{104} is H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, or (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het**, said alkyl, cycloalkyl,
 10 aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het** being optionally substituted with R^{150} ;
 - c) $OCOR^{105}$ wherein R^{105} is (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het**, said alkyl, cycloalkyl, aryl,
 15 **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het** being optionally substituted with R^{150} ;
 - d) SR^{108} , $SO_2N(R^{108})_2$ or $SO_2N(R^{108})C(O)R^{108}$ wherein each R^{108} is independently H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl or (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het** or both R^{108} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het** or heterocycle being optionally substituted with R^{150} ;
 - e) $NR^{111}R^{112}$ wherein R^{111} is H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl or (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het**, and R^{112} is H, CN, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl or (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl, (C_{1-6}) alkyl)**Het**, $COOR^{115}$ or SO_2R^{115} wherein R^{115} is (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, or (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het**, or both R^{111} and R^{112} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het**, or heterocycle being optionally substituted with R^{150} ;
 - f) $NR^{116}COR^{117}$ wherein R^{116} and R^{117} is each H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het**, said (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl,

23

- Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being optionally substituted with R¹⁵⁰;
- g) NR¹¹⁸CONR¹¹⁹R¹²⁰, wherein R¹¹⁸, R¹¹⁹ and R¹²⁰ is each H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, or R¹¹⁸ is covalently bonded to R¹¹⁹ and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle; or R¹¹⁹ and R¹²⁰ are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle;
- said alkyl, cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** or heterocycle being optionally substituted with R¹⁵⁰;
- h) NR¹²¹COCOR¹²² wherein R¹²¹ and R¹²² is each H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, a 6- or 10-membered aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being optionally substituted with R¹⁵⁰; or R¹²² is OR¹²³ or N(R¹²⁴)₂ wherein R¹²³ and each R¹²⁴ is independently H, (C₁₋₆alkyl), (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, or R¹²⁴ is OH or O(C₁₋₆alkyl) or both R¹²⁴ are covalently bonded together to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** and heterocycle being optionally substituted with R¹⁵⁰;
- i) COR¹²⁷ wherein R¹²⁷ is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said alkyl, cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being optionally substituted with R¹⁵⁰;
- j) COOR¹²⁸ wherein R¹²⁸ is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl and (C₁₋₆alkyl)**Het** being optionally substituted with R¹⁵⁰;
- k) CONR¹²⁹R¹³⁰ wherein R¹²⁹ and R¹³⁰ are independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, or both R¹²⁹ and R¹³⁰ are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl, (C₁₋₆alkyl)**Het** and heterocycle being optionally substituted with

R^{150} ;

l) aryl, Het, (C1-6alkyl)aryl or (C1-6alkyl)Het, all of which being optionally substituted with R^{150} ;

wherein R^{150} is preferably:

5 - 1 to 3 substituents selected from: halogen, NO_2 , cyano or azido;
or

- 1 to 3 substituents selected from:

10 **a)** (C₁₋₆) alkyl or haloalkyl, (C₃₋₇)cycloalkyl, (C₂₋₆)alkenyl, (C₂₋₈)alkynyl, (C₁₋₆) alkyl-(C₃₋₇)cycloalkyl, all of which optionally substituted with R^{160} ;

b) OR^{104} wherein R^{104} is H, (C₁₋₆)alkyl) or (C₃₋₇)cycloalkyl, said alkyl or cycloalkyl optionally substituted with R^{160} ;

15 **d)** SR^{108} , $SO_2N(R^{108})_2$ or $SO_2N(R^{108})C(O)R^{108}$ wherein each R^{108} is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, Het, or both R^{108} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, aryl, Het and heterocycle being optionally substituted with R^{160} ;

20 **e)** $NR^{111}R^{112}$ wherein R^{111} is H, (C₁₋₆)alkyl, or (C₃₋₇)cycloalkyl, and R^{112} is H, (C₁₋₆)alkyl or (C₃₋₇)cycloalkyl, $COOR^{115}$ or SO_2R^{115} wherein R^{115} is (C₁₋₆)alkyl or (C₃₋₇)cycloalkyl, or both R^{111} and R^{112} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl and heterocycle being optionally substituted with R^{160} ;

25 **f)** $NR^{116}COR^{117}$ wherein R^{116} and R^{117} is each H, (C₁₋₆)alkyl or (C₃₋₇)cycloalkyl said (C₁₋₆)alkyl and (C₃₋₇)cycloalkyl being optionally substituted with R^{160} ;

30 **g)** $NR^{118}CONR^{119}R^{120}$, wherein R^{118} , R^{119} and R^{120} is each H, (C₁₋₆)alkyl or (C₃₋₇)cycloalkyl, or R^{118} is covalently bonded to R^{119} and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle;
or R^{119} and R^{120} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated

25

heterocycle;

said alkyl, cycloalkyl, and heterocycle being optionally substituted with R^{160} ;

5

h) $NR^{121}COCOR^{122}$ wherein R^{121} is H, (C_{1-6}) alkyl or (C_{3-7}) cycloalkyl, said alkyl and cycloalkyl being optionally substituted with R^{160} ;

10

or R^{122} is OR^{123} or $N(R^{124})_2$ wherein R^{123} and each R^{124} is independently H, (C_{1-6}) alkyl or (C_{3-7}) cycloalkyl, or both R^{124} are covalently bonded together to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl and heterocycle being optionally substituted with R^{160} ;

15

i) COR^{127} wherein R^{127} is H, (C_{1-6}) alkyl or (C_{3-7}) cycloalkyl, said alkyl and cycloalkyl being optionally substituted with R^{160} ;

j) $COOR^{128}$ wherein R^{128} is H, (C_{1-6}) alkyl or (C_{3-7}) cycloalkyl, said (C_{1-6}) alkyl and (C_{3-7}) cycloalkyl being optionally substituted with R^{160} ; and

20

k) $CONR^{129}R^{130}$ wherein R^{129} and R^{130} are independently H, (C_{1-6}) alkyl or (C_{3-7}) cycloalkyl, or both R^{129} and R^{130} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl and heterocycle being optionally substituted with R^{160} ;

25

wherein R^{160} is defined as 1 or 2 substituents selected from:

halogen, CN, C_{1-6} alkyl, haloalkyl, $COOR^{161}$, OR^{161} , $N(R^{162})_2$, $SO_2N(R^{162})_2$, $NR^{162}COR^{162}$ or $CON(R^{162})_2$, wherein R^{161} and each R^{162} is independently H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl or (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl; or both R^{162} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle.

30


More preferably, R^2 is selected from: aryl or Het, each optionally monosubstituted or disubstituted with substituents selected from the group consisting of: halogen, haloalkyl, N_3 , or

a) (C_{1-6}) alkyl optionally substituted with OH, $O(C_{1-6})$ alkyl or $SO_2(C_{1-6})$

- alkyl);
- b) (C₁₋₆)alkoxy;
- e) $\text{NR}^{111}\text{R}^{112}$ wherein both R^{111} and R^{112} are independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or R^{112} is 6- or 10-membered aryl, **Het**, (C₁₋₆)alkyl-aryl or (C₁₋₆)alkyl-**Het**; or both R^{111} and R^{112} are covalently bonded together and to the nitrogen to which they are attached to form a nitrogen-containing heterocycle, each of said alkyl, cycloalkyl, aryl, **Het**, alkyl-aryl or alkyl-**Het**; being optionally substituted with halogen or:
- OR^{161} or $\text{N}(\text{R}^{162})_2$, wherein R^{161} and each R^{162} is independently H, (C₁₋₆)alkyl, or both R^{162} are covalently bonded together and to the nitrogen to which they are attached to form a nitrogen-containing heterocycle;
- f) NHCOR^{117} wherein R^{117} is (C₁₋₆)alkyl, O(C₁₋₆)alkyl or O(C₃₋₇)cycloalkyl;
- i) CO-aryl; and
- k) CONH_2 , $\text{CONH}(\text{C}_{1-6}\text{alkyl})$, $\text{CON}(\text{C}_{1-6}\text{alkyl})_2$, CONH-aryl , or $\text{CONHC}_{1-6}\text{alkyl-aryl}$.
- Still, more preferably, R^2 is aryl or **Het**, each optionally monosubstituted or disubstituted with substituents selected from the group consisting of: halogen, haloalkyl, or
- a) (C₁₋₆)alkyl optionally substituted with OH, O(C₁₋₆)alkyl or $\text{SO}_2(\text{C}_{1-6}\text{alkyl})$;
- b) (C₁₋₆)alkoxy; and
- e) $\text{NR}^{111}\text{R}^{112}$ wherein both R^{111} and R^{112} are independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or R^{112} is 6- or 10-membered aryl, **Het**, (C₁₋₆)alkyl-aryl or (C₁₋₆)alkyl-**Het**; or both R^{111} and R^{112} are covalently bonded together and to the nitrogen to which they are attached to form a nitrogen-containing heterocycle, each of said alkyl, cycloalkyl, aryl, **Het**, alkyl-aryl or alkyl-**Het**; or being optionally substituted with halogen or:
- OR^{161} or $\text{N}(\text{R}^{162})_2$, wherein R^{161} and each R^{162} is independently H, (C₁₋₆)alkyl, or both R^{162} are covalently bonded together and to the nitrogen to which they are


attached to form a nitrogen-containing heterocycle.

Even more preferably, R^2 is phenyl or a heterocycle selected from:




Even more preferably, R^2 is selected from the group consisting of:


H, Br, $CH=CH_2$,




28



29




Still more preferably, R^2 is selected from:



10

Most preferably, R^2 is selected from:



5

R^3 :

Preferably, R^3 is selected from (C_{3-7}) cycloalkyl, (C_{3-7}) cycloalkenyl, (C_{6-10}) bicycloalkyl, (C_{6-10}) bicycloalkenyl, 6- or 10-membered aryl, or Het. More preferably, R^3 is (C_{3-7}) cycloalkyl. Most preferably, R^3 is cyclopentyl, or cyclohexyl.

10

Y :

Preferably Y^1 is O.

Z :

- 15 Preferably, Z is OR^6 , wherein R^6 is H, (C_{1-6}) alkyl being optionally substituted with: halo, hydroxy, carboxy, amino, C_{1-6} alkoxy, C_{1-6} alkoxycarbonyl, and C_{1-6} alkylamino; or R^6 is C_{1-6} alkylaryl optionally substituted with: halogen, cyano, nitro, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkanoyl, $-(CH_2)_{1-6}-COOR^7$, $-(CH_2)_{1-6}-CONR^7R^8$, $-(CH_2)_{1-6}-NR^7R^8$, $-(CH_2)_{1-6}-NR^7COR^8$, $-(CH_2)_{1-6}-NH-SO_2R^7$, $-(CH_2)_{1-6}-OR^7$, $-(CH_2)_{1-6}-SR^7$, $-(CH_2)_{1-6}-SO_2R^7$, and $-(CH_2)_{1-6}-SO_2NR^7R^8$, wherein each R^7 and each R^8 is H or C_{1-6} alkyl,

- or Z is NR^9R^{10} wherein each of R^9 and R^{10} is selected from: H, C_{1-6} alkoxy, or C_{1-6} alkyl optionally substituted with halo, hydroxy, carboxy, amino, C_{1-6} alkoxy, C_{1-6} alkoxycarbonyl, and C_{1-6} alkylamino;

25

More preferably, Z is OH or O(C₁₋₆alkyl) or Z is NR⁹R¹⁰ wherein R⁹ is preferably H and R¹⁰ is preferably H or C₁₋₆alkyl.

5 Most preferably, Z is OH.

Specific embodiments

Included within the scope of this invention are all compounds of formula I as presented in Tables 1 and 2.

10

Polymerase activity

The ability of the compounds of formula (I) to inhibit RNA synthesis by the RNA dependent RNA polymerase of HCV can be demonstrated by any assay capable of measuring RNA dependent RNA polymerase activity. A suitable assay is described in the examples.

15

Specificity for RNA dependent RNA polymerase activity

To demonstrate that the compounds of the invention act by specific inhibition of HCV polymerase, the compounds may be tested for inhibitory activity in a DNA dependent RNA polymerase assay.

20

When a compound of formula (I), or one of its therapeutically acceptable salts, is employed as an antiviral agent, it is administered orally, topically or systemically to mammals, e.g. humans, rabbits or mice, in a vehicle comprising one or more pharmaceutically acceptable carriers, the proportion of which is determined by the solubility and chemical nature of the compound, chosen route of administration and standard biological practice.

25

For oral administration, the compound or a therapeutically acceptable salt thereof can be formulated in unit dosage forms such as capsules or tablets each containing a predetermined amount of the active ingredient, ranging from about 25 to 500 mg, in a pharmaceutically acceptable carrier.

30

For topical administration, the compound can be formulated in pharmaceutically accepted vehicles containing 0.1 to 5 percent, preferably 0.5 to 5 percent, of the

35

active agent. Such formulations can be in the form of a solution, cream or lotion.

For parenteral administration, the compound of formula (I) is administered by either intravenous, subcutaneous or intramuscular injection, in compositions with pharmaceutically acceptable vehicles or carriers. For administration by injection, it is preferred to use the compounds in solution in a sterile aqueous vehicle which may also contain other solutes such as buffers or preservatives as well as sufficient quantities of pharmaceutically acceptable salts or of glucose to make the solution isotonic.

10

Suitable vehicles or carriers for the above noted formulations are described in pharmaceutical texts, e.g. in "Remington's The Science and Practice of Pharmacy", 19th ed., Mack Publishing Company, Easton, Penn., 1995, or in "Pharmaceutical Dosage Forms And Drugs Delivery Systems", 6th ed., H.C. Ansel et al., Eds., Williams & Wilkins, Baltimore, Maryland, 1995.

The dosage of the compound will vary with the form of administration and the particular active agent chosen. Furthermore, it will vary with the particular host under treatment. Generally, treatment is initiated with small increments until the optimum effect under the circumstance is reached. In general, the compound of formula I is most desirably administered at a concentration level that will generally afford antivirally effective results without causing any harmful or deleterious side effects.

For oral administration, the compound or a therapeutically acceptable salt is administered in the range of 10 to 200 mg per kilogram of body weight per day, with a preferred range of 25 to 150 mg per kilogram.

For systemic administration, the compound of formula (I) is administered at a dosage of 10 mg to 150 mg per kilogram of body weight per day, although the aforementioned variations will occur. A dosage level that is in the range of from about 10 mg to 100 mg per kilogram of body weight per day is most desirably employed in order to achieve effective results.

When the compositions of this invention comprise a combination of a compound

of formula I and one or more additional therapeutic or prophylactic agent, both the compound and the additional agent should be present at dosage levels of between about 10 to 100%, and more preferably between about 10 and 80% of the dosage normally administered in a monotherapy regimen.


- 5 When these compounds or their pharmaceutically acceptable salts are formulated together with a pharmaceutically acceptable carrier, the resulting composition may be administered *in vivo* to mammals, such as man, to inhibit HCV polymerase or to treat or prevent HCV virus infection. Such treatment may also be achieved using the compounds of this invention in combination with agents
- 10 which include, but are not limited to: immunomodulatory agents, such as α -, β -, or γ -interferons; other antiviral agents such as ribavirin, amantadine; other inhibitors of HCV NS5B polymerase; inhibitors of other targets in the HCV life cycle, which include but not limited to, helicase, NS2/3 protease, NS3 protease, or internal ribosome entry site (IRES); or combinations thereof. The additional
- 15 agents may be combined with the compounds of this invention to create a single dosage form. Alternatively these additional agents may be separately administered to a mammal as part of a multiple dosage form.

Methodology and Synthesis

- 20 Indole derivatives or analogs according to the present invention can be prepared from known monocyclic aromatic compounds by adapting known literature sequences such as those described by J.W. Ellingboe et al. (*Tet. Lett.* **1997**, *38*, 7963) and S. Cacchi et al. (*Tet. Lett.* **1992**, *33*, 3915). Scheme 1, shown below wherein R^1 , R^2 , R^3 , R^6 , K, L, and M are as described herein illustrate how these
- 25 procedures can be adapted to the synthesis of compounds of formula 1 of this invention.

Scheme 1

34



In carrying out the route illustrated in Scheme 1, a suitably protected form of 3-trifluoroacetamido-4-iodobenzoic acid **I(i)** is reacted with an alkyne **I(ii)** in the presence of a metal catalyst (e.g. a palladium metal complex such as $\text{PdCl}_2(\text{PPh}_3)_2$, Pd_2dba_3 , $\text{Pd}(\text{PPh}_3)_4$ and the like), a base (Et_3N , DIEA and the like or an inorganic basic salt including metal carbonates, fluorides and phosphates), and optionally in the presence of an additional phosphine ligand (triaryl or heteroarylphosphine, dppe, dppf, dppp and the like). Suitable solvents for this reaction include DMF, dioxane, THF, DME, toluene, MeCN, DMA and the like at temperatures ranging from 20 °C to 170 °C, or alternatively without solvent by heating the components together. Alternatively, the cross-coupling reaction can be carried out on a suitably protected form of 3-amino-4-iodobenzoate and the amino group can be trifluoroacetylated in the subsequent step as described by J.W. Ellingboe et al. (*Tet. Lett.* **1997**, *38*, 7963).

Reaction of the above diarylalkynes **I(iii)** with an enol triflate under cross-coupling conditions similar to those described above gives after hydrogenation of the double bond, indole derivatives **I(iv)**. Enol triflates are known and can be prepared from the corresponding ketones by following known literature methods (for example, cyclohexene triflate can be prepared from cyclohexanone, triflic anhydride and a hindered organic base such as 2,6-di-*tert*-butyl-4-methylpyridine). The hydrogenation of the double bond originally present in R^1 can be carried out

35


with hydrogen gas or a hydrogen donor (ammonium formate, formic acid and the like) in the presence of a metal catalyst (preferably Pd) in a suitable solvent (lower alkyl alcohols, THF etc.).

- 5 Finally, following hydrolysis of the ester protecting group in **I(iv)**, the resulting 6-carboxyindole derivative **I(v)** is converted to compounds of formula **1** by coupling with the appropriate amine of formula H_2N-R^6 . Condensation of the 6-indolecarboxylic acid with amines H_2N-R^6 can be accomplished using standard amide bond forming reagents such as TBTU, HATU, BOP, BroP, EDAC, DCC,
- 10 isobutyl chloroformate and the like, or by activation of the carboxyl group by conversion to the corresponding acid chloride prior to condensation with an amine. Any remaining protecting group is removed following this step to give compounds of formula **1**.
- 15 Alternatively, compounds of formula **1** can be prepared by elaboration from a pre-existing indole core by following adaptations of literature procedures as described, for example, by P. Gharagozloo et al. (*Tetrahedron* **1996**, *52*, 10185) or K. Freter (*J. Org. Chem.* **1975**, *40*, 2525). Such a methodology is illustrated in Scheme 2;

20

Scheme 2

36



- In carrying the route illustrated in Scheme 2, commercially available 6-indolecarboxylic acid **2(i)**, which can also be prepared according to the method of S. Kamiya et al. (*Chem. Pharm. Bull.* **1995**, *43*, 1692) is used as the starting material. The indole **2(i)** is reacted with a ketone **2(ii)** under basic or acidic aldol-type conditions. Suitable conditions to affect this condensation include strong bases such as alkali metal hydroxides, alkoxides and hydrides in solvents such as lower alkyl alcohols (MeOH, EtOH, *tert*BuOH etc.), THF, dioxane, DMF, DMSO, DMA and the like at reaction temperature ranging from $-20^\circ C$ to $120^\circ C$. Alternatively, the condensation can be carried out under acid conditions using organic or mineral acids or both. Appropriate conditions include mixtures of AcOH and aqueous phosphoric acid at temperatures ranging from $15^\circ C$ to $120^\circ C$.
- Following protection of the carboxylic acid group in the form of an ester (usually lower alkyl) using known methods, the indole nitrogen can be alkylated with R^3 if desired. Reaction conditions to alkylate the nitrogen of an indole derivative are well known to those skilled in the art and include the use of strong bases such as alkali metal hydrides, hydroxides, amides, alkoxides and alkylmetals, in the appropriate solvent (such as THF, dioxane, DME, DMF, MeCN, DMSO, alcohols

37

and the like) at temperatures ranging from $-78\text{ }^{\circ}\text{C}$ to $140\text{ }^{\circ}\text{C}$. An electrophilic form of R^3 is used for the alkylation of the indole anion. Such electrophilic species include iodides, bromides, chlorides and sulfonate esters (mesylates, tosylate, brosylate or triflate).

- 5 Halogenation (usually bromination, but also iodination) of the 2-position of the indole **2(iv)** gives **2(v)**. Suitable halogenating agents include, for example, elemental bromine, *N*-bromosuccinimide, pyridine tribromide, dibromohydantoin and the corresponding iodo derivatives. Suitable solvents for this reaction are inert to reactive halogenating agents and include for example hydrocarbons,
- 10 chlorinated hydrocarbons (DCM , CCl_4 , CHCl_3), ethers (THF, DME, dioxane), acetic acid, ethyl acetate, IPA, and mixtures of these solvents. Reaction temperature ranges from $-40\text{ }^{\circ}\text{C}$ to $100\text{ }^{\circ}\text{C}$. A method of choice to carry out the bromination of indoles as shown in Scheme 2 was described by L. Chu (*Tet. Lett.* **1997**, *38*, 3871).

15


- The 2-bromoindole derivatives **2(v)** can be converted directly to fully substituted key intermediates **I(v)** through a cross-coupling reaction with aryl or heteroaryl boronic acids, boronate esters or trialkylstannane derivatives. These boron or tin organometallic species are from commercial sources or can be prepared by
- 20 standard literature procedures. Cross-coupling with organoboron reagents can be carried out by any variations of the Suzuki cross-coupling reaction reported in the literature. This usually involves the use of a transition metal catalyst (usually Pd^0), triaryl or triheteroarylphosphine ligands, an additive such as an inorganic chloride (e.g. LiCl), and a base (usually an aqueous inorganic base such as
- 25 sodium or potassium carbonate or phosphate). The reaction is usually carried out in an alcoholic solvent (EtOH), DME, toluene, THF and the like at temperatures ranging from $25\text{ }^{\circ}\text{C}$ to $140\text{ }^{\circ}\text{C}$.

- Cross-coupling with tin reagents can be carried out by any variations of the Stille cross-coupling reaction reported in the literature. This usually involves the use of
- 30 a transition metal catalyst (usually Pd^0), triaryl or triheteroaryl phosphine ligands, and an additive such as an inorganic chloride (e.g. LiCl) or iodide (e.g. CuI). Suitable solvents for this reaction include toluene, DMF, THF, DME and the like at temperatures ranging from $25\text{ }^{\circ}\text{C}$ to $140\text{ }^{\circ}\text{C}$. Intermediate **I(v)** is then converted to compounds of formula **1** as described for Scheme 1.

- 35 Alternatively, the 2-bromoindole intermediate **2(v)** can be trans-metallated to an

organotin species (or organozinc) and used in Stille-type cross-coupling reactions under conditions described above. In this case, aromatic and heteroaromatic halides (chlorides, bromides, iodides) or triflates are used to introduce R^2 . The conversion of 2-bromoindole derivatives **2(v)** to the corresponding organotin species **2(vi)** is carried out via initial low-temperature (usually -78° to -30°C) halogen-metal exchange using an alkyl lithium reagent (e.g. $n\text{BuLi}$ or $tert\text{-BuLi}$) or using lithium metal. The transient 2-lithioindole species is then trapped with a trialkyltin halide (e.g. $n\text{Bu}_3\text{SnCl}$ or Me_3SnCl). Alternatively, the lithioindole intermediate can be trapped with zinc chloride to form the corresponding organozincate which can also undergo transition metal-catalyzed cross-coupling with aromatic and heteroaromatic halides or triflates as described, for example, by M. Rowley (*J. Med. Chem.* **2001**, *44*, 1603).

The present invention also encompasses compounds of formula 1 where the carboxylic group is in the 5-position of the indole system. The synthesis of such compounds is based on adaptation of literature procedures and is depicted in Scheme 3:



20

In carrying out the synthetic route illustrated in Scheme 3, ethyl 4-acetamido-3-iodobenzoate **3(i)** undergoes metal catalyzed cross-coupling with an alkyne **3(ii)** to give a 2,3-disubstituted-5-indolecarboxylate **3(iii)** according to an adaptation of a procedure described by A. Bedeschi et al. (*Tet. Lett.* **1997**, *38*, 2307). The indole derivative **3(iii)** is then alkylated on nitrogen with electrophilic R^1 groups

25

(halides, sulfonate esters) under the action of a base such as alkali metal hydroxides, fluorides, hydrides amides, alkyllithium, phosphabases and the like, to give **3(iv)**. Suitable solvents for this alkylation include DMF, DMA, DMSO, MeCN, THF, dioxane, DME and the like. Following saponification of the ester group with an alkaline solution, the resulting 5-indolecarboxylic acid derivative **3(v)** is coupled to H₂N-R⁶ using an amide bond forming reagent as described previously (Scheme 1), to give compounds of formula I.

EXAMPLES

10


The present invention is illustrated in further detail by the following non-limiting examples. All reactions were performed in a nitrogen or argon atmosphere. Temperatures are given in degrees Celsius. Flash chromatography was performed on silica gel. Solution percentages or ratios express a volume to volume relationship, unless stated otherwise. Mass spectral analyses were recorded using electrospray mass spectrometry. Abbreviations or symbols used herein include:

- DIEA: diisopropylethylamine;
- DMAP: 4-(dimethylamino)pyridine;
- 20 DMSO: dimethylsulfoxide;
- DMF: N,N-dimethylformamide;
- Et: ethyl;
- EtOAc: ethyl acetate;
- Et₂O: diethyl ether;
- 25 HPLC: high performance liquid chromatography;
- Pr: isopropyl
- Me: methyl;
- MeOH: Methanol;
- MeCN: acetonitrile;
- 30 Ph: phenyl;
- TBE: tris-borate-EDTA;
- TBTU: 2-(1*H*-benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate;
- TFA: trifluoroacetic acid;
- TFAA: trifluoroacetic anhydride;
- 35 THF: tetrahydrofuran;

- MS (ES): electrospray mass spectrometry;
PFU: plaque forming units;
DEPC: diethyl pyrocarbonate;
DTT: dithiothreitol
- 5 EDTA: ethylenediaminetetraacetate
HATU: O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium
hexafluorophosphate
BOP: benzotriazole-1-yloxy-tris(dimethylamino)phosphonium
hexafluorophosphate
- 10 EDAC: see ECD
DCC: 1,3-Dicyclohexyl carbodiimide
HOBt: 1-Hydroxybenzotriazole
ES⁺: electro spray (positive ionization)
ES⁻: electro spray (negative ionization)
- 15 DCM: dichloromethane
TBME: *tert*-butylmethyl ether
TLC: thin layer chromatography
AcOH: acetic acid
EtOH: ethanol
- 20 DBU: 1,8-diazabicyclo[5.4.0]undec-7-ene
BOC: *tert*-butoxycarbonyl
Cbz: carbobenzyloxy carbonyl
*i*PrOH: isopropanol
NMP: N-methylpyrrolidone
- 25 EDC: 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride
RNasin: A ribonuclease inhibitor marketed by Promega Corporation
Tris: 2-amino-2-hydroxymethyl-1,3-propanediol
UMP: uridine 5'-monophosphate
UTP: uridine 5'-triphosphate
- 30 IPA: isopropyl acetate

Examples 1-22 illustrate methods of synthesis of representative compounds of this invention.

EXAMPLE 1

5 ***Methyl 3-amino-4-iodobenzoate:***

3-Amino-4-iodobenzoic acid (13.35 g, 50.8 mmol) was added to MeOH (150mL) and SOCl₂ (4.8 mL, 65.8 mmol, 1.3 equivalent) was added. The mixture was refluxed for 3 h and then volatiles were removed under reduced pressure. The residue was co-evaporated three times with MeOH and dried in vacuo (15.23 g).

10

Methyl 3-trifluoroacetamido-4-iodobenzoate:

The aniline derivative from above (14.53 g, 52 mmol) was dissolved in DCM (200 mL) and TFAA (15 mL, 104 mmol) was added. The dark purple solution was refluxed overnight. Volatiles were removed under reduced pressure and the residue was passed through a short pad of silica gel using DCM as eluent. The desired product was obtained as a pink solid (13.81 g).

15

4-Phenylethynyl-3-(2,2,2-trifluoro-ethanoylamino)-benzoic acid methyl ester:

The iodide from above (0.742 g, 2 mmol), phenylacetylene (0.37 mL, 3.9 mmol, 1.7 equivalent) and Et₃N (6 mL) were charged in a dry flask under argon.

20 PdCl₂(PPh₃)₂ (0.241 g, 0.3 mmol) was added and the mixture was stirred at room temperature until judged complete by HPLC analysis (~5 h). The reaction mixture

was concentrated to half volume under reduced pressure and diluted with water (80 mL). The mixture was extracted with EtOAc (3 x 100 mL) and the organic extract washed with 5% HCl (100 mL), after (100 mL) and brine (40 mL). After drying over MgSO₄, the residue was purified by flash chromatography using 20% EtOAc – hexane as eluent to give the desired cross-coupled alkyne as a tan solid (0.442 g).

Methyl 3-(cyclohexenyl)-2-phenylindole 6-carboxylate:

A flame-dried flask was charged with finely powdered anhydrous K₂CO₃ (0.153 g, 1.1 mmol) and the alkyne derivative from above (0.390 g, 1.1 mmol). Dry DMF (4 mL) was added and the suspension degassed with a stream of argon. The enol triflate derived from cyclohexanone, prepared following the procedure described by A.G. Martinez, M. Hanack et al. (*J. Heterocyclic Chem.* **1988**, *25*, 1237 or equivalent methods described in the literature, (0.802 g, 3.3 mmol, 3 equivalents) was added followed by Pd(PPh₃)₄ (0.086 g, 0.07 mmol) and the mixture was stirred for 8 h at room temperature. DMF was removed under vacuum and the residue purified by flash chromatography using DCM as eluent (0.260 g).

Methyl 3-cyclohexyl-2-phenylindole-6-carboxylate:

The material from above was hydrogenated (1 atm H₂ gas) over 20% Pd(OH)₂ in the usual manner, using MeOH as solvent. The desired cyclohexane indole was isolated after filtration of the catalyst.


3-Cyclohexyl-2-phenylindole-6-carboxylic acid:

The methyl ester from above (0.154 g, 0.15 mmol) was refluxed overnight in a mixture of MeOH (10 mL) and 2N NaOH (6 mL) until complete hydrolysis had occurred as shown by HPLC analysis. After cooling to room temperature, 2N HCl (5 mL) was added followed by AcOH to pH 7. MeOH was removed under reduced pressure, water (50 mL) was added and the product extracted with EtOAc. The extract was washed with water and brine, and dried (MgSO₄). Removal of volatiles under reduced pressure gave the title indole carboxylic acid as a light-orange solid (0.149 g).

Following the same procedure but using 2-ethynylpyridine instead of phenylacetylene, 3-cyclohexane-2-(2-pyridyl)indole-6-carboxylic acid was

obtained.

EXAMPLE 2:



5

3-Cyclohexenyl-6-indole carboxylic acid:

A 12 L round-bottomed flask was equipped with a reflux condenser and a mechanical stirrer, and the system was purged with nitrogen gas. 6-Indole carboxylic acid (300.00 g, 1.86 mole, 3 equivalents) was charged into the flask, followed by MeOH (5.5 L). After stirring for 10 min at room temperature, cyclohexanone (579 mL, 5.58 mole) was added. Methanolic sodium methoxide (25% w/w, 2.6 L, 11.37 mole, 6.1 equivalents) was added in portions over 10 min. The mixture was then refluxed for 48 h. After cooling to room temperature, water (4 L) was added and methanol removed under reduced pressure. The residual aqueous phase was acidified to pH 1 with concentrated HCl (~1.2 L). The resulting yellowish precipitate was collected by filtration, washed with water and dried under vacuum at 50 °C. The desired cyclohexane derivative was obtained as a beige solid (451.0g, 100% yield).

3-Cyclohexyl-6-indole carboxylic acid:

The unsaturated derivative from above was hydrogenated for 20 h under 55 psi hydrogen gas pressure over 20% Pd(OH)₂/C (10.25 g) using 1:1 THF – MeOH (2.5 L) as solvent. After filtration of the catalyst, volatiles were removed under reduced pressure and the residue was triturated with hexane. The beige solid was collected by filtration, washed with hexane and dried under vacuum (356.4 g, 78% yield).

Methyl 3-cyclohexyl-6-indole carboxylate:

A 5 L three-necked flask was equipped with a reflux condenser and a mechanical stirrer, and the system was purged with nitrogen gas. The indole carboxylic acid from above (300.00 g, 1.233 mole) was charged into the flask and suspended in MeOH (2 L). Thionyl chloride (5 mL, 0.0685 mole, 0.05 equivalent) was added dropwise and the mixture was refluxed for 48 h. Volatiles were removed under reduced pressure and the residue was triturated with hexane to give a beige solid that was washed with hexane and dried under vacuum (279.6 g, 88% yield).

Methyl-2-bromo-3-cyclohexyl-6-indole carboxylate:


Adapting the procedure of L. Chu (*Tet. Lett.* **1997**, *38*, 3871) methyl 3-cyclohexyl-6-indole carboxylate (4.65 g, 18.07 mmol) was dissolved in a mixture of THF (80 mL) and CHCl₃ (80 mL). The solution was cooled in an ice bath and pyridinium bromide perbromide (pyridine tribromide, 7.22 g, 22.6 mmol, 1.25 equivalent) was added. After stirring for 1.5 h at 0 °C, the reaction was judged complete by TLC. It was diluted with CHCl₃ (200 mL), washed with 1M NaHSO₃ (2 x 50 mL), saturated aqueous NaHCO₃ (2 x 50 mL) and brine (50 mL). After drying over Na₂SO₄, the solvent was removed under reduced pressure and the residue crystallized from TBME – hexane. The desired 2-bromoindole derivative was collected by filtration, washed with hexane and dried (3.45 g). Evaporation of mother liquors gave a red solid that was purified by flash chromatography using 15% EtOAc in hexane yielding an additional 3.62 g of pure material. Total yield was 5.17 g (85% yield).

EXAMPLE 3:***General procedure for the Suzuki cross-coupling of aryl and heteroarylboronic acids with 2-bromoindole derivatives:***

Cross-coupling of aromatic/heteroaromatic boronic acid or ester derivatives with 2-bromoindoles such as the one described in example 2 can be performed using any variations of the standard metal-catalyzed Suzuki cross-coupling reaction as described in the literature and well known to those skilled in the art. The following example serves to illustrate such a process and is non-limiting.

3-Cyclohexyl-2-furan-3-yl-1H-indole-6-carboxylic acid methyl ester:


45



The 2-bromoindole of example 2 (8.92 g, 26.5 mmol), 3-furanboronic acid (B.P. Roques et al. *J. Heterocycl. Chem.* **1975**, *12*, 195; 4.45 g, 39.79 mmol, 1.5 equivalent) and LiCl (2.25 g, 53 mmol, 2 equivalents) were dissolved in a mixture of EtOH (100 mL) and toluene (100 mL). A 1M aqueous Na₂CO₃ solution (66 mL, 66 mmol) was added and the mixture was degassed with argon for 45 min. Pd(PPh₃)₄ (3.06 g, 2.65 mmol, 0.1 equivalent) was added and the mixture stirred overnight at 75-85 °C under argon. Volatiles were removed under reduced pressure and the residue re-dissolved in EtOAc (500 mL). The solution was washed with water, saturated NaHCO₃ (100 mL) and brine (100 mL). After drying over a mixture of MgSO₄ and decolorizing charcoal, the mixture was filtered and concentrated under reduced pressure. The residual oil was triturated with a mixture of TBME (20 mL) and hexane (40 mL), cooled in ice and the precipitated solid collected by filtration, washed with cold 25% TBME in hexane, and dried (3.09 g). The filtrate and washings from the above trituration were combined, concentrated and purified by flash chromatography using 10-25% EtOAc in hexane to give an additional 4.36 g of product. The total yield of the 2-(3-furyl)indole of example 3 was 8.25 g.

EXAMPLE 4:

46



Methyl 3-cyclohexyl-1-methyl-6-indole carboxylate:

Methyl 3-cyclohexyl-6-indole carboxylate from example 2 (150.00 g, 0.583 mole) was charged into a 3 L three-necked flask equipped with a mechanical stirrer and purged with nitrogen gas. DMF (1 L) was added and the solution was cooled in an ice-bath. NaH (60% oil dispersion, 30.35 g, 0.759 mole, 1.3 equivalent) was added in small portions (15 min) and the mixture was stirred for 1 h in the cold. Iodomethane (54.5 mL, 0.876 mole, 1.5 equivalent) was added in small portions, maintaining an internal temperature between 5 – 10 °C. The reaction mixture was then stirred overnight at room temperature. The reaction was quenched by pouring into ice-water (3 L), resulting in the formation of a cream-colored precipitate. The material was collected by filtration, washed with water and dried in vacuum at 45 °C (137.3 g, 86% yield).

Methyl 2-bromo-3-cyclohexyl-1-methyl-6-indole carboxylate:


The 1-methylindole derivative from above (136.40 g, 0.503 mole) was charged into a 5 L three-necked flask equipped with a mechanical stirrer and purged with nitrogen gas. CHCl₃ (750 mL) and THF (750 mL) were added and the solution was cooled to 0 °C. Pyridine tribromide (pyridinium bromide perbromide, 185.13 g, 0.579 mole, 1.15 equivalent) was added in small portions and the mixture was stirred for 1 h at 0 °C. The solvent was removed under reduced pressure at room temperature and the residue dissolved in EtOAc (3 L). The solution was washed with water and brine, dried (decolourising charcoal / MgSO₄) and concentrated under reduced pressure. The residue was suspended in TBME and heated to 50 °C. The suspension was stored overnight in the refrigerator and the cream-

47

coloured crystalline product was collected by filtration. It was washed with TBME and dried in vacuum (134.3 g, 76% yield).

EXAMPLE 5:***Cyclohexyl-methyl-tributylstannanyl-1H-indole-6-carboxylic acid methyl ester:***

5



The bromoindole derivative of example 4 (2.70 g, 7.71 mmol) was dissolved in dry THF (40 mL) and the solution was cooled to -78°C under an argon atmosphere.

10 A solution of $n\text{BuLi}$ in hexane (1.4 M, 6.90 mL, 9.64 mmol, 1.25 equivalent) was added dropwise over 15 min and stirring at low temperature was continued for 75 min. To the resulting suspension was added $n\text{Bu}_3\text{SnCl}$ (2.93 mL, 10.8 mmol, 1.4 equivalent) over 5 min. The suspension dissolved and the solution was stirred for 1 h at -78°C . The reaction mixture was warmed to room temperature and THF


15 removed under reduced pressure. The residue was dissolved in TBME (150 mL), washed with 1:1 brine – water and dried over MgSO_4 . The material was purified by chromatography on silica gel that was previously deactivated by mixing with a solution of 5% Et_3N in hexane. The same solvent was used as eluent for the chromatography. The title stannane was isolated as a yellow oil (3.42 g, 79 %

20 yield).

EXAMPLE 6:**General procedure for Stille cross-coupling of the 2-stannane indole of example 5 with aromatic/heteroaromatic halides:**

25 Cross-coupling of aromatic/heteroaromatic halides or pseudohalides (preferably bromides, iodides and triflates) with the stannane derivative of example 5 can be performed using any variations of the standard metal-catalyzed Stille cross-coupling reaction as described in the literature. The following example serves to illustrate such a process.

3-Cyclohexyl-1-methyl-2-pyridin-2-yl-1H-indole-6-carboxylic acid methyl ester:




5

The stannane derivative of example 5 (3.42 g, 6.1 mmol) was dissolved in DMF (10 mL) and CuI (0.116 g, 0.61 mmol, 0.1 equivalent), LiCl (0.517 g, 12.21 mmol, 2 equivalent), triphenylphosphine (0.320 g, 1.22 mmol, 0.2 equivalent) and 2-bromopyridine (0.757 mL, 7.94 mmol, 1.3 equivalent) were added. The solution was degassed with a stream of argon (30 min) and Pd(PPh₃)₄ (0.352 g, 0.31 mmol, 0.05 equivalent) was added. After purging with argon for an additional 10 min, the solution was heated and stirred at 100 °C overnight under argon. The DMF was then removed under vacuum and the residue dissolved in EtOAc (150 mL). The solution was washed with 1N NaOH (25 mL) and brine (25 mL) and dried over MgSO₄. The solvent was removed under reduced pressure and the residue purified by flash chromatography eluting with CHCl₃ then 5-10% EtOAc in CHCl₃ (1.516 g, 71% yield).

EXAMPLE 7:

General procedure for Stille cross-coupling of 2-bromoindoles with aryl or heteroaryl stannanes:

3-Cyclohexyl-1-methyl-2-pyridin-2-yl-1H-indole-6-carboxylic acid methyl ester:



The 2-bromoindole derivative of example 4 (0.150 g, 0.428 mmol) and 2-

49

trimethylstannylthiophene (S.F. Thames et al. J. Organometal. Chem. 1972, 38, 29; 0.150 g, 0.61 mmol, 1.4 equivalent) were dissolved in dry THF (7 mL) in a sealed tube, and the solution was degassed with a stream of argon for 30 min. Pd(Cl)₂(PPh₃)₂ (0.018 g, 0.026 mmol, 0.06 equivalent) was added and the tube
5 sealed. The solution was heated to 80 °C for 40 h. The reaction mixture was cooled to room temperature, EtOAc (10 mL) was added and the suspension filtered. After evaporation of the solvent, the residue was re-submitted to the reaction conditions for an additional 20 h, with fresh 2-stannylthiophene (0.150 g, 0.61 mmol) and catalyst (0.020 g). After cooling to room temperature and
10 filtration of solids, the solvent was evaporated and the residue purified by flash chromatography using 15-100% CHCl₃ in hexane as eluent (0.133 g, 88% yield).

The same procedure can be used to couple stannane derivatives to the 2-bromoindole of Example 2.


15

EXAMPLE 8:

General procedure for the N-alkylation of 2-aryl and 2-heteroaryl-6-indole carboxylates:

3-Cyclohexyl-1-methyl-2-pyridin-2-yl-1H-indole-6-carboxylic acid methyl ester:

20



NaH (60% oil dispersion, 0.186 g, 4.64 mmol, 1.5 equivalent) was washed with hexane (20 mL) to remove the oil and then re-suspended in DMF (5 mL). After cooling to 0 °C in an ice bath, the indole derivative of example 3 (1.000 g, 3.09
25 mmol) was added dropwise as a solution in DMF (3 mL + 2 mL rinse). After stirring for 15 min, iodomethane (0.385 mL, 6.18 mmol, 2 equivalents) was added in one portion and the mixture was stirred for 2 h in the cold and an additional 2 h at room temperature. The reaction was then quenched by addition of 1N HCl (1 mL) and diluted with TBME (100 mL). The solution was washed with 1N HCl (25
30 mL) and dried (MgSO₄). After removal of volatiles under reduced pressure, the residue was purified by flash chromatography using 5-10% EtOAc in hexane as

eluent to give the title compound as a white solid (0.903 g, 86% yield).

Other *N*-alkylindole derivatives within the scope of this invention could be prepared from the appropriate electrophiles (e.g. EtI, iPrI, iBuI, BnBr) using a similar procedure.

5


EXAMPLE 9:

General procedure for the saponification of 6-indolecarboxylates to the corresponding free carboxylic acids:

This procedure applies to both indole and *N*-methylindole carboxylates.

10

3-Cyclohexyl-1-methyl-2-pyridin-2-yl-1H-indole-6-carboxylic acid:




- 15 The 6-indole carboxylate of example 6 (1.517 g, 4.35 mmol) was dissolved in DMSO (8 mL) and 5N NaOH (4.4 mL) was added. The mixture was stirred at 50 °C for 30 min. The solution was then cooled to room temperature and added dropwise to water (15 mL). Insoluble black impurities were removed by filtration and AcOH (2 mL) was added dropwise to the filtrate. The white precipitate that
- 20 formed was collected by filtration, washed with water and dried (1.37 g, 94% yield).

EXAMPLE 10:

1-Cyclohexyl-2-phenyl-1H-indole-5-carboxylic acid:

51

**Ethyl 4-amino-3-iodobenzoate:**

Ethyl 4-aminobenzoate (15.00 g, 91 mmol) and iodine (11.80 g, 46.5 mmol) were mixed with water (80 mL) and chlorobenzene (4.60 g, 41 mmol). The mixture was stirred while the temperature was gradually raised to 90 °C over 30 min. Hydrogen peroxide (30%, 50 mL) was added over 10 h at 90 °C. After stirring at that temperature for an additional 6 h, the mixture was cooled and the solution decanted from the residual solids. The solids were dissolved in DCM and the solution washed successively with sodium thiosulfate and brine. After drying (MgSO₄), the solvent was removed under reduced pressure and the resulting brown solid was triturated with hexane to remove di-iodinated by-products. The desired compound was obtained as a brown solid (22.85 g, 86% yield).

Ethyl 4-acetamido-3-iodobenzoate:

The aniline from above (1.00 g, 3.44 mmol) was dissolved in pyridine (5 mL) and the solution was cooled in ice. AcCl (0.32 mL, 4.47 mmol, 1.3 equivalent) was added dropwise and the mixture was stirred for 1 h at 0 °C and 2 h at room temperature. The reaction mixture was then diluted with 1 N HCl and the product was extracted with TBME (100 mL). The organic phase was washed with 1N HCl (50 mL), dried (MgSO₄) and concentrated to give the desired material as a tan-colored solid (1.121 g, 97% yield).

Ethyl 2-phenyl-indole-5-carboxylate:

Following the procedure of A. Bedeschi et al. (*Tet. Lett.* **1997**, *38*, 2307), the acetanilide derivative from above (0.900 g, 2.7 mmol) was reacted with phenylacetylene (0.385 mL, 3.5 mmol, 1.3 equivalent) in the presence of

52

PdCl₂(PPh₃)₂ (10 mole %) and CuI (10 mole %) in a mixture of dioxane (5 mL) and tetramethylguanidine (5 mL). The desired 2-phenylindole (0.589 g, 82% yield) was isolated as a yellow solid after flash chromatography with 15% EtOAc in hexane.

5

1-Cyclohex-1-enyl-2-phenyl-1H-indole-5-carboxylic acid ethyl ester:

The 2-phenylindole derivative from above (0.265 g, 1.0 mmol) was dissolved in DMF (2 mL) and cesium hydroxide monohydrate (0.208 g, 1.2 mmol, 1.2 equivalent) was added. The solution was cooled in an ice bath and 3-bromocyclohexene (0.193 g, 1.2 mmol, 1.2 equivalent) was added dropwise (5 min) as a solution in DMF (1 mL). The mixture was stirred for 30 min at 0 °C. The reaction was diluted with water (25 mL), extracted with Et₂O (2 x 50 mL) and the extract dried over MgSO₄. The solvent was evaporated under reduced pressure to give a white foam (0.095 g) that was used without purification in the next step.

15


1-Cyclohexyl-2-phenyl-1H-indole-5-carboxylic acid:

The crude indole from above was hydrogenated in the usual way (1 atm H₂ gas) in EtOH over 20% Pd(OH)₂ on carbon for 20 h at room temperature. After filtration of the catalyst, the EtOH was removed under reduced pressure. The residue was dissolved in a mixture of MeOH (1 mL) and DMSO (1 mL) and 5N NaOH (0.5 mL) was added. The mixture was stirred overnight at 50 °C. The reaction mixture was cooled and water (10 mL) was added. After acidification with 1N HCl, the product was extracted into Et₂O (70 mL) and the solution dried (Na₂SO₄). Evaporation of the solvent gave a green residue consisting of a 2:1 mixture (85 mg) of the desired 1-cyclohexyl-2-phenylindole-5-carboxylic acid and 1,3-dicyclohexyl-2-phenylindole-5-carboxylic acid.

25

EXAMPLE 11:***1-Cyclohexyl-3-methyl-2-phenyl-1H-indole-5-carboxylic acid:***

53



Ethyl 2-phenyl-3-methyl-indole-5-carboxylate:

Adapting the procedure of H.-C. Zhang (*Tet. Lett.* **1997**, *38*, 2439) ethyl 4-amino-3-iodobenzoate (from example 10, 0.500 g, 1.72 mmol) was dissolved in DMF (5 mL) and LiCl (0.073 g, 1.72 mmol, 1 equivalent), PPh₃ (0.090 g, 0.34 mmol, 0.2 equivalent), K₂CO₃ (1.188 g, 8.6 mmol, 5 equivalents) and phenylpropyne (0.645 mL, 5.76 mmol, 3 equivalents) were added. The solution was degassed by purging with argon for 1 h and palladium acetate (0.039 g, 0.17 mmol, 0.1 equivalent) was added. The mixture was stirred at 80 °C under argon for 20 h. The reaction mixture was diluted with water (25 mL) and extracted with EtOAc (50 mL). The extract was washed with brine (3 x 25 mL) and dried (MgSO₄). Concentration under reduced pressure and purification by flash chromatography with 10-15% EtOAc – hexane gave the desired 2-phenyl-3-methyl indole (0.275 g, least polar component) and the 3-phenyl-2-methyl isomer (0.109 g, more polar component).

Ethyl 1-(3-cyclohexenyl)-3-methyl-2-phenylindole-5-carboxylate:

The less polar isomer from above (0.264 g, 0.95 mmol) was dissolved in DMSO (2 mL) and cesium hydroxide monohydrate (0.191 g, 1.14 mmol, 1.2 equivalent) was added followed by 3-bromocyclohexene (0.183 g, 1.14 mmol, 1.2 equivalent in 0.7 mL of DMSO). The mixture was stirred at room temperature for 30 min. Additional CsOH monohydrate (0.400 g, 2.4 equivalents) and 3-bromocyclohexene (0.400 g, 2.4 equivalents) were added and stirring continued for an additional 30 min. Similar amounts of the two reagents were again added and after another 30 min of stirring at room temperature, the reaction was diluted with 1N HCl (6 mL) and water (20 mL). The product was extracted with TBME

(100 mL), dried (MgSO_4) and after concentration under reduced pressure, the residue was purified by flash chromatography using 5-10% EtOAc in hexane as eluent. The desired *N*-alkylated indole was obtained (0.130 g).

5 **Ethyl 1-cyclohexyl-3-methyl-2-phenylindole-5-carboxylate:**


The unsaturated product from above was hydrogenated (1 atm H_2 gas) in the usual way over 20% $\text{Pd}(\text{OH})_2$ in EtOH at room temperature for 3 h.

1-Cyclohexyl-3-methyl-2-phenyl-1H-indole-5-carboxylic acid:

10 The hydrogenated material from above was dissolved in a mixture of DMSO (2 mL) and MeOH (2 mL). 5N NaOH (0.5 mL) was added and the mixture was stirred overnight at 60 °C. After dilution with water (40 mL), the product aqueous phase was washed with a 1:1 mixture of Et_2O – hexane (50 mL) and then acidified with 1N HCl to pH 1. The liberated free acid was extracted with diethyl ether (2 x 50 mL) and the extract dried over Na_2SO_4 . Removal of the solvent
15 under reduced pressure gave the desired indole as a light brown solid (0.074 g).

EXAMPLE 12:

2-Bromo-3-cyclopentyl-1-methyl-1H-indole-6-carboxylic acid methyl ester:



20 A 3 L three-necked flask equipped with a mechanical stirrer was charged with indole 6-carboxylic acid (220 g, 1.365 mole) and KOH pellets (764.45 g, 13.65 mole, 10 equivalents). Water (660 mL) and MeOH (660 mL) were added and the mixture heated to 75 °C. Cyclopentanone (603.7 mL, 6.825 mole, 5 equivalents) was added dropwise over 18 h using a pump. The reaction mixture was heated for an additional 3 h (after which the reaction was judged complete by HPLC) and
25 cooled to 0 °C for 1 h. The precipitated potassium salt is collected by filtration, and washed with TBME (2 X 500 mL) to remove cyclopentanone self-condensation products. The brown solid was re-dissolved in water (2.5 L) and the

55

solution washed with TBME (2 X 1 L). Following acidification to pH 3 with conc. HCl (425 mL), the beige precipitate was collected by filtration, washed with water (2 X 1 L) and dried under vacuum at 70 °C. The crude product weighed 275.9 g (88.9 % mass recovery) and had an homogeneity of 85% (HPLC).

5

The crude product from above (159.56 g, 0.70 mole) was dissolved in MeOH (750 mL) and 20% Pd(OH)₂ on charcoal (8.00 g) was added. The mixture was hydrogenated in a Parr apparatus under 50 psi hydrogen gas for 18 h. After completion, the catalyst was removed by filtration through celite and the solvent removed under reduced pressure. The resulting brown solid was dried at 70 °C under vacuum for 12 h. The crude product (153.2 g) was obtained as a brown solid and was 77% homogeneous by HPLC.

The crude 3-cyclopentylindole-6-carboxylic acid (74.00 g, 0.323 mole) was charged in a 3 L three-necked flask equipped with a mechanical stirrer and a thermometer. The system was purged with nitrogen gas and anhydrous DMF (740 mL) was added. After dissolution of the starting material, anhydrous potassium carbonate (66.91 g, 0.484 mole, 1.5 equivalent) was added and the mixture stirred for 5 minutes. Iodomethane (50 mL, 0.807 mole, 2.5 equivalents) was added and the mixture stirred for 5 h after which HPLC analysis of the reaction mixture indicated 97% conversion to the methyl ester.

The reaction mixture was cooled in an ice bath and sodium hydride (95%, oil-free, 10.10 g, 0.420 mole, 1.3 equivalent) was added in small portions over 3 min (exothermic: 8 °C to 30 °C internal temperature raise). After stirring for an additional 15 min, the cooling bath was removed and stirring continued for 1.5 h at room temperature after which no further progression was observed (HPLC). Additional NaH (1.55 g, 65 mmol, 0.2 equivalent) and iodomethane (1.0 mL, 16 mmol, 0.05 equivalent) were added and after stirring for 15 min, the reaction was judged complete by HPLC (96% N-methylated).


The reaction mixture was slowly (2 min) poured into water (4 L) with vigorous stirring and after 10 min, acidified to pH <2 with conc. HCl (85 mL). The mixture was stirred for 5 min to allow complete conversion of any remaining potassium carbonate and bicarbonate to the more soluble chloride. The pH was adjusted to ~7 with 4N NaOH (40 mL) and the mixture stirred overnight at room temperature. The precipitated material was collected by filtration, washed with water (600 mL)

56

and dried at 60 °C under vacuum. The crude product (79% homogeneity by HPLC) was obtained as a brown solid (72.9 g).

The crude material from above is triturated with a minimal amount of MeOH to remove a series of minor impurities. The solid was then collected by filtration and dissolved in a minimal amount of hot EtOAc. After cooling to room temperature, hexane was added (5 X volume) and the mixture cooled in ice and filtered. The filtrate was then evaporated to dryness to give the desired product.

The *N*-methylindole from above (10.60 g, 41.2 mmol) was dissolved in isopropyl acetate (150 mL) and sodium acetate (5.07 g, 62 mmol, 1.5 equivalent) was added. The suspension was cooled in an ice bath and bromine (2.217 mL, 43.3 mmol, 1.05 equivalent) was added dropwise over 2 min. The pale amber suspension turned dark red (exotherm from 5 °C to 13 °C). It was stirred for 1 h at 0 °C. The reaction was completed by adding additional bromine (0.21 mL, 4.2 mmol, 0.10 equivalent) as shown by HPLC analysis. The reaction was then quenched by addition of 10% aqueous sodium sulfite solution (15 mL), followed by water (50 mL) and K₂CO₃ (10.6 g, 1.8 equivalent) to neutralize HBr. The organic layer was separated, washed with 10% aqueous sodium sulfite and aqueous K₂CO₃ and dried (MgSO₄). The solvent was removed under reduced pressure and the residue co-evaporated with TBME (75 mL) to give a beige solid that was dried under vacuum overnight (13.80 g). The crude material was triturated with boiling MeOH (80 mL) for 30 min, cooled in ice and the beige solid collected by filtration. The product was dried at 60 °C under vacuum (10.53 g, 76% recovery).

EXAMPLE 13***3-Cyclopentyl-1-methyl-2-vinyl-1H-indole-6-carboxylic acid:***


To the 2-bromoindole derivative of example 12 (2.044 g, 6.08 mmol) in dry dioxane (20 mL) was added vinyltributyltin (1.954 mL, 6.69 mmol). The solution was degassed by bubbling nitrogen for 15 min. Then bis(triphenylphosphine)

palladium (II) chloride (213.4 mg, 0.304 mmol) was added and the reaction mixture was heated at 100 °C overnight. The reaction mixture was diluted with ether and successively washed with water and brine. After the usual treatment (MgSO₄, filtration and concentration) the residue was flash chromatographed (5
5 cm, 10% AcOEt-hexane) to afford the desired compound (1.32 g, 4.70 mmol, 77 % yield) as a white solid.

To the ester from above (153 mg, 0.54 mmol) in a mixture of THF (2.8 mL) and methanol (1.4 mL) was added an aqueous solution of lithium hydroxide (226.6 mg, 5.40 mmol in 1.6 mL of water). The reaction mixture was stirred at 50 °C for
10 1.5 h and diluted with water. The aqueous layer was acidified with 1M aq. HCl and extracted three times with CH₂Cl₂. The combined organic layers were successively washed with water (2X) and brine. After the usual treatment (MgSO₄, filtration and concentration) the desired crude acid was isolated (150 mg).

15 EXAMPLE 14

3-Cyclohexyl-1-methyl-2-oxazol-5-yl-1H-indole-6-carboxylic acid:



To the bromide of example 4 (1.00 g, 2.855 mmol) in dry dioxane (10 mL) was added vinyltributyltin (917.8 µL, 3.141 mmol). The solution was degassed by
20 bubbling nitrogen through for 15 min. Then bis(triphenylphosphine) palladium (II) chloride (101 mg, 0.144 mmol) was added and the solution was refluxed for 7 hrs. The reaction mixture was diluted with ether and successively washed with water and brine. After the usual treatment (MgSO₄, filtration and concentration) the residue was flash chromatographed (5 cm, hexane to 2.5% AcOEt to 5% AcOEt
25 to 10% AcOEt-hexane) to afford the desired compound (773 mg, 2.60 mmol, 91 % yield) as a pale yellow solid.

To the olefinic derivative from above (100 mg, 0.336 mmol) in a mixture of

58

acetone (690 μ L), *tert*-butanol (690 μ L) and water (690 μ L) were successively added *N*-methylmorpholine *N*-oxide (NMMO; 48 mg, 0.410 mmol) and a 2.5 % solution of osmium tetroxide in *tert*-butanol (33 μ L). The reaction mixture was stirred at room temperature for three days and then concentrated. The residue
5 was dissolved in EtOAc and successively washed with water (2X) and brine. After the usual treatment (MgSO_4 , filtration and concentration) the crude diol (117 mg) was isolated.

To the crude diol obtained above (ca. 0.336 mmol) in a mixture of THF (3.2 mL)
10 and water (3.2 mL) at 0 °C was added sodium periodate (86.2 mg, 0.403 mmol). The cooling bath was then removed and the reaction mixture was stirred at room temperature for 1h 45 min. AcOEt was then added. The resulting solution was successively washed with 10% aq. citric acid, water, satd aq. NaHCO_3 , water (2X) and brine. After the usual treatment (MgSO_4 , filtration and concentration) the
15 crude desired aldehyde was isolated (92 mg, 0.307 mmol, 91 % yield).


A mixture of the aldehyde from above (25.8 mg, 0.086 mmol), anhydrous potassium carbonate (12.4 mg, 0.090 mmol) and Tosmic (17.57 mg, 0.090 mmol) in absolute MeOH (500 μ L) was refluxed for 2 h. AcOEt was then added and the
20 mixture was successively washed with water (2X) and brine. After the usual treatment (MgSO_4 , filtration and concentration) the crude desired oxazole was isolated (28 mg, 0.083 mmol, 96 % yield).

To the ester from above (28 mg, 0.083 mmol) in a mixture of THF (425 μ L),
25 MeOH (210 μ L) and water (250 μ L) was added lithium hydroxide (34.8 mg, 0.830 mmol). The reaction mixture was stirred overnight at room temperature, then diluted with water and acidified with a 1N aq. HCl solution. The aqueous layer was extracted with dichloromethane (3X) and successively washed with water (2X) and brine. After the usual treatment (MgSO_4 , filtration and concentration) the title
30 crude acid was isolated (30 mg).

EXAMPLE 15


2-(1H-Benzimidazol-2-yl)-3-cyclohexyl-1-methyl-1H-indole-6-carboxylic acid:

59



To a mixture of the aldehyde from example 14 (28 mg, 0.094 mmol) and 1,2-diaminobenzene (10.9 mg, 0.101 mmol) in acetonitrile (500 μ L) and DMF (200 μ L) was added chloranil (24.8 mg, 0.101 mmol). The reaction mixture was stirred at room temperature for three days. AcOEt was added and the reaction mixture was successively washed with a 1N aq. NaOH solution (2X), water (4X) and brine. After the usual treatment (MgSO₄, filtration and concentration) the residue was flash chromatographed (1 cm, 30% AcOEt-hexane) to afford the desired benzimidazole ester derivative (11 mg, 0.028 mmol, 30 % yield).

To the ester from above (11 mg, 0.028 mmol) in a mixture of THF (240 μ L), MeOH (120 μ L) and water (140 μ L) was added lithium hydroxide (11.7 mg, 0.280 mmol). The reaction mixture was stirred overnight at room temperature, then diluted with water and acidified with a 1N aq. HCl solution. The aqueous layer was extracted with dichloromethane (3X) and successively washed with water (2X) and brine. After the usual treatment (MgSO₄, filtration and concentration) the title crude acid was isolated (9 mg, 0.0241 mmol, 86 % yield).


EXAMPLE 16***3-Cyclopentyl-1-methyl-1H-indole-2,6-dicarboxylic acid 6-methyl ester:***

To the 3-cyclopentyl aldehyde prepared in a similar fashion to that described in example 15 (20 mg, 0.07 mmol) and 2-methyl-2-butene (541 μ L, 5.11 mmol) in tert-butanol (500 μ L) at 0 °C was added a freshly prepared solution of sodium chlorite (64.2 mg, 0.711 mmol) in phosphate buffer (98 mg of NaH₂PO₄ in 150 μ L of water). The reaction mixture was stirred for 45 min. at room temperature then brine was added. The aqueous layer was extracted twice with EtOAc. The

combined organic layer was successively washed with a 0.5 N aq. HCl solution and brine. After the usual treatment (MgSO₄, filtration and concentration) 23.1 mg of the desired crude acid were isolated as a yellow solid.

5 EXAMPLE 18

3-Cyclopentyl-2-pyridin-2-yl-benzofuran-6-carboxylic acid:




- The 2-bromobenzofuran derivative of example 17 (0.850 g, 2.93 mmol), 2-tri(n-butyl)stannylpyridine (1.362 g, 3.7 mmol), triphenylphosphine (0.760 g, 2.90 mmol), lithium chloride (0.250 g, 5.9 mmol) and CuI (0.057 g, 0.3 mmol) were dissolved in DMF (30 mL) and the mixture was degassed by bubbling argon for 30 min. Tetrakis (triphenylphosphine)palladium (0.208 g, 0.18 mmol) was added and the mixture stirred at 100 °C under an argon atmosphere. After 19 h, the reaction was cooled to room temperature, poured into water (70 mL) and extracted with TBME. The organic phase was washed with water (2 X) and brine, dried (MgSO₄) and concentrated to give a residue that was purified by flash chromatography. The desired 2(2-pyridyl)benzofuran derivative (0.536 g, 63 % yield) was obtained as a white solid.
- The nitrile from above (0.200 g, 0.694 mmol) was suspended in a mixture of conc. H₂SO₄ (5 mL), AcOH (4 mL) and water (2 mL). After refluxing for 1.5 h, TLC showed complete hydrolysis. The mixture was cooled in ice and the 10 N NaOH was added dropwise to pH 9. The aqueous layer was washed with dichloromethane and then acidified to pH 6 with 5 N HCl. The product was extracted with EtOAc, dried (MgSO₄) and solvents removed under reduced pressure. The desired carboxylic acid was obtained as a white solid.

EXAMPLE 19

2-Bromo-3-cyclopentyl-benzo[b]thiophene-6-carboxylic acid ethyl ester

61



To a solution of 3-bromo-6-cyclopentanecarbonylphenol of Example 17 (5.194 g, 19.30 mmol) in DMF (58.0 mL) was added 1,4-diazabicyclo[2.2.2]octane (4.33 g, 38.60 mmol) and dimethylthiocarbamyl chloride (4.77 g, 38.6 mmol) at room temperature. The mixture was stirred at room temperature for 3 hr. The mixture was acidified with 1 N HCl to pH 3 and then extracted with EtOAc. The organic layers were combined and washed with brine and dried over MgSO₄. The crude mixture was purified through a plug of silica gel with 3%EtOAc/hexanes to provide 6.976 g (100%) of the desired thiocarbamate as a colorless oil.

- 5 The neat *O*-3-bromo-6-cyclopentanecarbonyl *N,N*-dimethylthiocarbamate from above (43.147 g, 121.1 mmol) was heated to internal temperature of 180-190 °C for 5 hr. TLC (20% EtOAc/hexanes: *R*_f 0.6 (starting material), 0.5 (product)) was used to monitor the reaction progress. The crude material was used for the next reaction without further purification.
- 15 The crude *S*-3-bromo-6-cyclopentanecarbonyl *N,N*-dimethylthiocarbamate from above was dissolved in MeOH (600 mL), KOH (40.0 g, 714 mmol) was added and the mixture was heated to reflux for 1.5 h. The mixture was cooled to room temperature and the solvent was removed by rotary evaporation. The residue was dissolved in water and acidified by 6 N HCl to pH 3. It was extracted with EtOAc and the crude product was purified by a silica gel chromatography with 1-5% EtOAc/hexanes. 31.3 g (91%) of the desired thiophenol derivative was obtained as a yellow oil.
- 20

To a solution of the 3-bromo-6-cyclopentanecarbonylthiophenol from above (0.314 g, 1.105 mmol) in acetone (5.0 mL) was added K₂CO₃ (0.477 g, 3.45

mmol) followed by addition of ethyl bromoacetate (0.221 g, 0.147 mL, 1.33 mmol).

The mixture was stirred overnight. The reaction mixture was filtered through filter paper and the filtrate was concentrated. Purification by silica gel with 5% EtOAc/hexanes provided 0.334 g (82%) of the product as a colorless oil.

5 The crude ester from above was dissolved in THF (12.0 mL), 1 N NaOH (5.0 mL) was added at room temperature. The mixture was stirred at room temperature for 2-3 hr, or until TLC indicated complete reaction. The solvent was removed by rotary evaporation. Water was added and the mixture was acidified with 6 N HCl to pH 3 and extracted with EtOAc, washed with brine and dried over MgSO₄. The
10 solvent was removed under reduced pressure and the residue was used without further purification.

To the crude acid from above was added acetic anhydride (16.0 mL), and then NaOAc (0.573 g) and the mixture was heated to reflux overnight. The mixture was cooled to room temperature and poured into a mixture of ice and toluene. 6
15 N NaOH was added until pH to about 7, and extracted with EtOAc, washed with brine and dried over MgSO₄. The solvent was removed by rotary evaporation and the residue was purified by silica gel with hexanes to provide 0.795 g (80%) of 6-bromo-3-cyclopentyl benzothiophene as a colorless oil.

A mixture of the 6-bromo-3-cyclopentylbenzothiophene from above (0.723 g, 2.57
20 mmol), and copper cyanide (0.272 g, 3.04 mmol) in DMF (1.4 mL) was heated to reflux overnight. The mixture was cooled to room temperature and diluted with EtOAc. 2 N NH₄OH was added and the mixture was stirred for 10 minutes and filtered through Celite. The aqueous layer was extracted with EtOAc. The organic layers were combined and washed with brine, dried over MgSO₄, and the
25 solvent was removed under reduced pressure. The product was used without further purification.


3-cyclopentyl-6-cyanobenzothiophene (17.65 g, 77.65 mmol) was dissolved in acetic acid (310 mL), bromine (49.64 g, 310.6 mmol) was added at room temperature. The mixture was stirred at room temperature overnight and HPLC
30 was used to monitor the reaction progress. After the reaction was complete, toluene was added to the reaction mixture to remove acetic acid (3 x 100 mL). The crude product was dried under reduced pressure and used without further purification.

The crude cyano derivative from above was added to ethanol (150 mL,
35 denatured) and conc. H₂SO₄ (45 mL) and the mixture heated to reflux for 1-2

days. After completion (HPLC) the reaction mixture was cooled to room temperature and poured into ice-water and extracted with dichloromethane (5 x 100 mL), the organic layers were combined and washed with 5% NaHCO₃, and brine. The solvent was removed under reduced pressure and the residue was
5 purified with silica gel by 1% EtOAc/hexanes. The collected fractions were concentrated and the residue was slurried in methanol. The solid was filtered and washed with ice-cold methanol to provide 15.9 g (58%, two steps) of pure ethyl ester as a slight yellow solid.

10 **EXAMPLE 20**

3-Cyclopentyl-2-pyridin-2-yl-benzo[b]thiophene-6-carboxylic acid:



The 2-bromobenzothiophene of example 19 (0.354 g, 1.00 mmol), 2-tri(n-butyl)stannylpyridine (0.442 g, 1.2 mmol), triphenylphosphine (0.262 g, 1.00
15 mmol), lithium chloride (0.085 g, 2.0 mmol) and CuI (0.019 g, 0.1 mmol) were dissolved in DMF (10 mL) and the mixture was degassed by bubbling argon for 30 min. Tetrakis (triphenylphosphine)palladium (0.069 g, 0.06 mmol) was added and the mixture stirred at 100 °C under an argon atmosphere. After 24 h, the reaction was cooled to room temperature, poured into water (70 mL) and extracted with
20 TBME. The organic phase was washed with water (2 X) and brine, dried (MgSO₄) and concentrated to give a residue that was purified by flash chromatography. The desired 2(2-pyridyl)benzothiophene ester (0.197 g, 56 % yield) was obtained as a pale yellow waxy solid.


The ester from above was hydrolyzed in the usual manner using NaOH, to give
25 the title acid that could be used directly or purified by HPLC and flash chromatography.

The acid could be coupled to amine derivatives following the general procedure described in example 37.


30 **EXAMPLE 21**

3-Cyclopentyl-2-furan-3-yl-benzo[b]thiophene-6-carboxylic acid:

64



The 2-bromobenzothiophene ester of example 19 was coupled to 3-furanboronic acid as described in example 3 to give the desired 2(3-furyl)benzothiophene ester in 85 % yield. Saponification of the ethyl ester was carried out with NaOH at room temperature to give the title carboxylic acid derivative.

EXAMPLE 22**3-Cyclohexyl-1-methyl-2-phenyl-1H-pyrrolo[2,3,b]pyridine-6-carboxylic acid:**

10

7-Azaindole (15.00 g, .127 mole) was dissolved in MeOH (330 mL) and sodium methoxide (25% w/w in MeOH, 172 mL, 0.753 mole) and cyclohexanone (52.86 mL, 0.51 mole) were added. The mixture was refluxed for 60 h and then concentrated under reduced pressure. After cooling in ice-water, the reaction mixture was acidified to pH 8 with 3N HCl and the precipitated solid was collected by filtration. The product was washed with water, triturated with TBME-hexane and dried by azeotroping with toluene (19.8 g).

The material from above (15.00 g, 75.65 mmol) was dissolved in a mixture of EtOH (130 mL) and THF (30 mL) and 20% Pd(OH)₂ on carbon (1.30 g) was added. The mixture was hydrogenated under 1 atm of H₂ gas for 24 h, after which point additional catalyst (1.30 g) was added. After stirring under H₂ gas for an additional 16 h, the catalyst was removed by filtration and the solution evaporated under reduced pressure to give a residue that was triturated with TBME to give an amber solid (13.9 g).

65

The azaindole derivative from above (7.50 g, 37.45 mmol) was dissolved in DME (130 mL) and *meta*-chloroperbenzoic acid (12.943 g, 60.0 mmol) was added. After stirring for 2 h, volatiles were removed under reduced pressure and the residue suspended in water (100 mL). The mixture was basified to pH 10 by
5 addition of saturated aqueous Na₂CO₃ solution under vigorous stirring. The solid was then collected by filtration, washed with water and a small amount of TBME, and dried (7.90 g).

The crude N-oxide from above (4.00 g, 18.49 mmol) was dissolved in DMF (350 mL) and NaH (60% dispersion, 1.52 g, 38 mmol) was added in small portions
10 over 5 min. The mixture was stirred for 30 min and iodomethane (1.183 mL, 19 mmol) was added dropwise over 20 min to the suspension. After stirring for 3 h at room temperature, no more progress was measured by HPLC analysis. The reaction mixture was poured into water and extracted 3 times with EtOAc. The extract was washed with brine, dried (MgSO₄) and evaporated to give an amber
15 solid (3.65 g, 60% homogeneity by NMR) that was used immediately without purification.

The crude product from above (0.80 g, 3.47 mmol) was dissolved in MeCN (10 mL). Triethylamine (1.13 mL, 8.1 mmol) was added followed by trimethylsilyl cyanide (2.13 mL, 16 mmol). The solution was then refluxed for 19 h. After
20 cooling to room temperature, the reaction was quenched by slow addition of aqueous NaHCO₃ and the product extracted with EtOAc. The extract was washed with brine, dried (MgSO₄) and concentrated to a residue that was purified by flash chromatography on silica gel using 15% EtOAc-hexane (0.285 g).

The nitrile (0.300 g, 1.254 mmol) was suspended in EtOH (15 mL) and hydrogen
25 chloride gas was bubbled through for 15 min to give a clear solution. The solution was then refluxed for 1.5 h until TLC showed complete conversion of starting material. After cooling to room temperature, volatiles were removed under reduced pressure and the residue was dissolved in EtOAc. The solution was washed with brine, dried (MgSO₄) and concentrated. The residue was purified by
30 flash chromatography on silica gel (15-20% EtOAc-hexane) to give the desired ethyl ester as a pale yellow gum (0.227 g).

The ester from above (0.100 g, 0.35 mmol) was dissolved in THF (4 mL) and pyridinium hydrobromide perbromide (0.200 g, 0.532 mmol) was added. The mixture was stirred at 65 °C in a sealed vial for 16 h (>80% conversion). The
35 solution was evaporated and the residue taken up into EtOAc. The solution was

washed with water and brine, dried (MgSO_4) and concentrated. The crude material was purified by flash chromatography on silica gel (15% EtOAc-hexane). The bromide from above (0.100 g, 0.274 mmol), phenylboronic acid (0.049 g, 0.4 mmol) and lithium chloride (0.019 g, 0.45 mmol) were dissolved in a mixture of
5 toluene (2 mL), EtOH (2 mL) and 1M Na_2CO_3 (0.43 mL). The mixture was degassed by passing argon gas through the solution for 30 min, and tetrakis(triphenylphosphine) palladium (0.035 g, 0.03 mmol) was added. The mixture was refluxed for 18 h after which point more catalyst (0.035 g, 0.03 mmol) was added. After refluxing for an additional 2 h, the EtOH was removed under
10 reduced pressure. The residue was dissolved in EtOAc and the solution washed with 10% aqueous HCl and brine, and dried (MgSO_4). Removal of volatiles under reduced pressure gave an orange gum that was purified by flash chromatography on silica gel using 20% EtOAc-hexane (0.105 g, crude).
The partially purified ester from above (0.100 g, 0.276 mmol) was dissolved in a
15 mixture of THF (2 mL) and EtOH (2 mL). 1N NaOH (2.8 mL) was added and the mixture stirred for 4 h at room temperature. Volatiles were removed under reduced pressure and the residue diluted with 10% aqueous HCl. The product was extracted with EtOAc (3 X), dried (MgSO_4), evaporated and purified by reversed-phase preparative HPLC to give the title compound.

20

EXAMPLE 23: INHIBITION OF NS5B RNA DEPENDENT RNA POLYMERASE ACTIVITY

The compounds of the invention were tested for inhibitory activity against the hepatitis C virus RNA dependant polymerase (NS5B), according to the following assay:

25 The substrates are:

a 12 nucleotide RNA oligo-uridylate (or oligo-uridine-monophosphate) (oligo-U) primer modified with biotin at the free 5'C position;

a complementary poly-adenylate (or adenosine monophosphate) (polyA) template of heterogeneous length (1000-10000 nucleotides); and

30 UTP-[5,6 ^3H].

Polymerase activity is measured as the incorporation of UMP-[5,6 ^3H] into the chain elongated from the oligo-U primer. The ^3H -labelled reaction product is captured by SPA-beads coated with streptavidin and quantified on the TopCount.

35 All solutions were made from DEPC treated MilliQ water [2 ml of DEPC is added

67

to 1 L of MilliQ water; the mixture is shaken vigorously to dissolve the DEPC, then autoclaved at 121°C for 30 minutes].

Enzyme: The full length HCV NS5B (**SEQ ID NO.1**) was purified as an N-terminal hexa-histidine fusion protein from baculovirus infected insect cells. The enzyme can be stored at -20°C in storage buffer (see below). Under these conditions, it was found to maintain activity for at least 6 months.

Substrates: The biotinylated oligo-U₁₂ primer, the Poly(A) template, and the UTP-[5,6 ³H] were dissolved in water. The solutions can be stored at -80°C.

10

Assay buffer:

20 mM Tris-HCl pH 7.5

5 mM MgCl₂

25 mM KCl

1 mM EDTA

15

1 mM DTT

NS5B storage buffer:

0.1 µM NS5B

25 mM Tris-HCl pH 7.5

300 mM NaCl

20

5 mM DTT

1 mM EDTA

0.1 % n-Dodecyl maltoside

30 % glycerol

Test compound cocktail: Just prior to assay, test compounds of the invention were dissolved in assay buffer containing 15% DMSO.

Substrate cocktail: Just prior to assay, the substrates were mixed in assay buffer to the following concentrations:

30

Component	Concentration in substrate cocktail	Final Concentration in assay
-----------	--	------------------------------------

68

RNAasin™	0.5 U/ μ L	1.67 U/ μ L
Biotin-oligo-U ₁₂ primer	3 ng/ μ L	1 ng/ μ L
PolyA template	30 ng/ μ L	10 ng/ μ L
UTP-[5,6- ³ H] 35 Ci/mmol	0.025 μ Ci/ μ L	0.0083 μ Ci/ μ L 0.25 μ M
UTP	2.25 μ M	0.75 μ M

Enzyme cocktail: Just prior to assay, the RNA polymerase (NS5B) cocktail was prepared in assay buffer to the following specifications:

Component	Concentration in cocktail
Tris-HCl at pH 7.5	20 mM
MgCl ₂	5 mM
KCl	25 mM
EDTA	1 mM
DTT	1 mM
n- Dodecyl maltoside	1%
NS5B	30 nM

5

Protocol:

The assay reaction was performed in a Microfluor™ white "U" bottom plate (Dynatech™ #7105), by successively adding:

20 μ L of test compound cocktail;

10 20 μ L of substrate cocktail; and

20 μ L of enzyme cocktail

(final [NS5B] in assay = 10 nM; final [n-dodecyl maltoside] in assay = 0.33%; final DMSO in assay = 5%).

The reaction was incubated at room temperature for 1.5 hours. STOP solution

15 (20 μ L; 0.5 M EDTA, 150 ng/ μ L tRNA) was added, followed by 30 μ L streptavidin coated PVT beads (8mg/ml in 20 mM Tris-HCl, pH 7.5, 25 mM KCl, 0.025% NaN₃). The plate was then shaken for 30 minutes. A solution of CsCl was added (70 μ L, 5 M), to bring the CsCl concentration to 1.95 M. The mixture was then allowed to stand for 1 hour. The beads were then counted on a Hewlett Packard

TopCount™ instrument using the following protocol:

Data mode: counts per minute

Scintillator: liq/plast

Energy range: low

5 Efficiency mode: normal

Region: 0-50

Count delay: 5 minutes

Count time: 1 minute

Expected results: 6000 cpm/well

10 200 cpm/well no enzyme control.

Based on the results at ten different concentrations of test compound, standard concentration-% inhibition curves were plotted and analysed to determine IC₅₀'s for the compounds of the invention. For some compounds the IC₅₀ was estimated from two points.

15

EXAMPLE 24: SPECIFICITY OF NS5B RNA DEPENDENT RNA POLYMERASE INHIBITION

The compounds of the invention were tested for inhibitory activity against polio virus RNA dependent RNA polymerase and calf thymus DNA dependent RNA polymerase II in the format that is described for the HCV polymerase with the exception that another polymerase was used in place of the HCV NS5B polymerase.

20

25 EXAMPLE 25: CELL BASED HCV RNA REPLICATION ASSAY

Cell Culture

Huh7 cells that stably maintain a subgenomic HCV replicon were established as previously described (Lohman et al., 1999. Science **285**: 110-113) and designated as the S22.3 cell-line. S22.3 cells are maintained in Dulbecco's Modified Earle Medium (DMEM) supplemented with 10% FBS and 1mg/mL neomycin (Standard Medium). During the assay, DMEM medium supplemented with 10% FBS, containing 0.5% DMSO and lacking neomycin was used (Assay Medium). 16 hours prior to compound addition, S22.3 cells are trypsinized and diluted to 50 000 cells/ml in Standard Medium. 200μL (10 000 cells) are

30

35

70

distributed into each well of a 96-well plate. The plate was then incubated at 37°C with 5% CO₂ until the next day.

Reagents and Materials:

5

Product	Company	Catalog #	Storage
DMEM	Wisent Inc.	10013CV	4°C
DMSO	Sigma	D-2650	RT
Dulbecco's PBS	Gibco-BRL	14190-136	RT
Fetal Bovine Serum	Bio-Whittaker	14-901F	-20°C/4°C
Neomycin (G418)	Gibco-BRL	10131-027	-20°C/4°C
Trypsin-EDTA	Gibco-BRL	25300-054	-20°C/4°C
96-well plates	Costar	3997	RT
PVDF 0.22µm Filter Unit	Millipore	SLGV025LS	RT
Deep-Well Titer Plate Polypropylene	Beckman	267007	RT

Preparation of Test Compound

10µL of test compound (in 100% DMSO) was added to 2 ml of Assay Medium for a final DMSO concentration of 0.5% and the solution was sonicated for 15 min and filtered through a 0.22µm Millipore Filter Unit. 900µl was transferred into row A of a Polypropylene Deep-Well Titer Plate. Rows B to H, contain 400µL aliquots of Assay Medium (containing 0.5% DMSO), and are used to prepare serial dilutions (1/2) by transferring 400µl from row to row (no compound was included in row H).

15 Application of test compound to cells

Cell culture medium was aspirated from the 96-well plate containing the S22.3 cells. 175µL of assay medium with the appropriate dilution of test compound was transferred from each well of the compound plate to the corresponding well of the cell culture plate (row H was used as the "No inhibition control"). The cell culture plate was incubated at 37°C with 5% CO₂ for 72 hours.

Extraction of Total Cellular RNA

Following the 72 hour incubation period, the total cellular RNA was extracted from the S22.3 cells of the 96-well plate using the RNeasy 96 kit (Qiagen®, RNeasy

71

Handbook. 1999.). Briefly, assay medium was completely removed from cells and 100 μ L of RLT buffer (Qiagen®) containing 143 mM β -mercaptoethanol was added to each well of the 96-well cell-culture plate. The microplate was gently shaken for 20 sec. 100 μ L of 70% ethanol was then added to each microplate well, and mixed by pipetting. The lysate was removed and applied to the wells of a RNeasy 96 (Qiagen®) plate that was placed on top of a Qiagen® Square-Well Block. The RNeasy 96 plate was sealed with tape and the Square-Well Block with the RNeasy 96 plate was loaded into the holder and placed in a rotor bucket of a 4K15C centrifuge. The sample was centrifuged at 6000 rpm (~5600 x g) for 4 min at room temperature. The tape was removed from the plate and 0.8 ml of Buffer RW1 (Qiagen® RNeasy 96 kit) was added to each well of the RNeasy 96 plate. The RNeasy 96 plate was sealed with a new piece of tape and centrifuged at 6000 rpm for 4 min at room temperature. The RNeasy 96 plate was placed on top of another clean Square-Well Block, the tape removed and 0.8 ml of Buffer RPE (Qiagen® RNeasy 96 kit) was added to each well of the RNeasy 96 plate. The RNeasy 96 plate was sealed with a new piece of tape and centrifuged at 6000 rpm for 4 min at room temperature. The tape was removed and another 0.8 ml of Buffer RPE (Qiagen® RNeasy 96 kit) was added to each well of the RNeasy 96 plate. The RNeasy 96 plate was sealed with a new piece of tape and centrifuged at 6000 rpm for 10 min at room temperature. Tape was removed, the RNeasy 96 plate was placed on top of a rack containing 1.2-mL collection microtubes. The RNA was eluted by adding 50 μ L of RNase-free water to each well, sealing plate with a new piece of tape and incubated for 1 min at room temperature. The plate was then centrifuged at 6000 rpm for 4 min at room temperature. The elution step was repeated with a second volume of 50 μ L RNase-free water. The microtubes with total cellular RNA are stored at -70°C.

Quantification of Total Cellular RNA

RNA was quantified on the STORM® system (Molecular Dynamics®) using the RiboGreen® RNA Quantification Kit (Molecular Probes®). Briefly, the RiboGreen reagent was diluted 200-fold in TE (10mM Tris-HCl pH =7.5, 1mM EDTA). Generally, 50 μ L of reagent was diluted in 10mL TE. A Standard Curve of ribosomal RNA was diluted in TE to 2 μ g/mL and pre-determined amounts (100, 50, 40, 20, 10, 5, 2 and 0 μ L) of the ribosomal RNA solution are then transferred in a new 96-well plate (COSTAR # 3997) and the volume was completed to 100 μ L

72

with TE. Generally, column 1 of the 96-well plate was used for the standard curve and the other wells are used for the RNA samples to be quantified. 10 μ L of each RNA sample that was to be quantified, was transferred to the corresponding well of the 96-well plate and 90 μ L of TE was added. One volume (100 μ L) of diluted

5 RiboGreen reagent was added to each well of the 96-well plate and incubated for 2 to 5 minutes at room temperature, protected from light (a 10 μ L RNA sample in a 200 μ L final volume generates a 20 X dilution). The fluorescence intensity of each well was measured on the STORM® system (Molecular Dynamics®). A standard curve was created on the basis of the known quantities of the ribosomal

10 RNA and the resulting fluorescent intensities. The RNA concentration in the experimental samples was determined from the standard curve and corrected for the 20X dilution.

Reagents and Materials:

15

Product	Company	Catalog #	Storage
DEPC	Sigma	D5758	4°C
EDTA	Sigma	E5134	RT
Trizma-Base	Sigma	T8524	RT
Trizma-HCl	Sigma	T7149	RT
Collection Tube Strips	Qiagen	19562	RT
Ribogreen RNA Quantitation Kit	Molecular Probe	R11490	-20°C
Rneasy 96 Kit	Qiagen	74183	RT
Square-Well Blocks	Qiagen	19573	RT

Real-Time RT-PCR

The Real-Time RT-PCR was performed on the ABI Prism 7700 Sequence Detection System using the TaqMan EZ RT-PCR Kit from (Perkin-Elmer Applied

20 Biosystems®). RT-PCR was optimized for the quantification of the 5' IRES of HCV RNA by using the Taqman technology (Roche Molecular Diagnostics Systems) similar to the technique previously described (Martell et al., 1999. J. Clin. Microbiol. 37: 327-332). The system exploits the 5'-3' nucleolytic activity of AmpliTaq DNA polymerase. Briefly, the method utilizes a dual-labeled fluorogenic

25 hybridization probe (PUTR Probe) that specifically anneals to the template

between the PCR primers (primers 8125 and 7028). The 5' end of the probe contains a fluorescent reporter (6-carboxyfluorescein [FAM]) and the 3' end contains a fluorescent quencher (6-carboxytetramethylrhodamine [TAMRA]). The FAM reporter's emission spectrum was suppressed by the quencher on the intact hybridization probe. Nuclease degradation of the hybridization probe releases the reporter, resulting in an increase in fluorescence emission. The ABI Prism 7700 sequence detector measures the increase in fluorescence emission continuously during the PCR amplification such that the amplified product was directly proportion to the signal. The amplification plot was analysed early in the reaction at a point that represents the logarithmic phase of product accumulation. A point representing a defined detection threshold of the increase in the fluorescent signal associated with the exponential growth of the PCR product for the sequence detector was defined as the cycle threshold (C_T). C_T values are inversely proportional to the quantity of input HCV RNA; such that under identical PCR conditions, the larger the starting concentration of HCV RNA, the lower the C_T . A standard curve was created automatically by the ABI Prism 7700 detection system by plotting the C_T against each standard dilution of known HCV RNA concentration.

Reference samples for the standard curve are included on each RT-PCR plate. HCV Replicon RNA was synthesized (by T7 transcription) *in vitro*, purified and quantified by OD₂₆₀. Considering that 1 µg of this RNA = 2.15×10^{11} RNA copies, dilutions are made in order to have 10^8 , 10^7 , 10^6 , 10^5 , 10^4 , 10^3 or 10^2 genomic RNA copies / 5 µL. Total cellular Huh-7 RNA was also incorporated with each dilution (50 ng / 5 µL). 5 µL of each reference standard (HCV Replicon + Huh-7 RNA) was combined with 45 µL of Reagent Mix, and used in the Real-Time RT-PCR reaction.

The Real-Time RT-PCR reaction was set-up for the experimental samples that were purified on RNeasy 96 –well plates by combining 5 µl of each total cellular RNA sample with 45 µL of Reagent Mix.

Reagents and Materials:

Product	Company	Catalog #	Storage
TaqMan EZ RT-PCR Kit	PE Applied Biosystems	N808-0236	-20°C
MicroAmp Optical Caps	PE Applied Biosystems	N801-0935	RT
MicroAmp Optical 96-	PE Applied Biosystems	N801-0560	RT

Well Reaction Plate			
---------------------	--	--	--

Reagent Mix preparation:

Component	Volume for one sample (μL)	Volume for One Plate (μL) (91 samples + Dead Volume)	Final conc.
Rnase-free water	16.5	1617	
5X TaqMan EZ buffer	10	980	1X
Mn(OAc) ₂ (25mM)	6	588	3mM
dATP (10mM)	1.5	147	300μM
dCTP (10mM)	1.5	147	300μM
dGTP (10mM)	1.5	147	300μM
dUTP (20mM)	1.5	147	600μM
Forward Primer (10μM)	1	98	200nM
Reverse Primer (10μM)	1	98	200nM
PUTR probe (5μM)	2	196	200nM
rTth DNA polymerase (2.5 U/μL)	2	196	0.1 U/μL
AmpErase UNG (1U/μL)	0.5	49	0.01 U/μL
Total Volume	45	4410	

Forward Primer Sequence (SEQ ID. 2): 5' - ACG CAG AAA GCG TCT AGC
 5 CAT GGC GTT AGT - 3'

Reverse Primer Sequence (SEQ ID NO. 3): 5' - TCC CGG GGC ACT CGC
 AAG CAC CCT ATC AGG - 3'

10 **Note:** Those primers amplify a region of 256-nt present within the 5' untranslated region of HCV.

PUTR Probe Sequence (SEQ ID NO. 4): 6FAM - TGG TCT GCG GAA CCG
 GTG AGT ACA CC - TAMRA

15

75

No Template Controls (NTC): On each plate, 4 wells are used as "NTC". For these controls, 5µl of water are added to the well in place of RNA.

Thermal Cycling Conditions:

5	50°C	2 min	} for 2 cycles
	60°C	30 min	
	95°C	5 min	
	95°C	15 sec	
10	60°C	1 min	} for 40 cycles
	90°C	15 sec	
	60°C	1 min	
	60°C	1 min	

Following the termination of the RT-PCR reaction the data analysis requires setting of threshold fluorescence signal for the PCR plate and a standard curve was constructed by plotting the Ct value versus RNA copy number used in each reference reaction. The Ct values obtained for the assay samples are used to interpolate an RNA copy number based on the standard curve.

Finally, the RNA copy number was normalized (based on the RiboGreen RNA quantification of the total RNA extracted from the cell culture well) and expressed as genome equivalents / µg of total RNA [ge/µg].

The RNA copy number [g.e./µg] from each well of the cell culture plate was a measure of the amount of replicating HCV RNA in the presence of various concentrations of inhibitor. The % inhibition was calculated with the following equation:

$$100 - [(g.e./\mu g \text{ inh}) / (g.e./\mu g \text{ ctl}) \times 100].$$

A non-linear curve fit with the Hill model was applied to the inhibition-concentration data, and the 50% effective concentration (EC₅₀) was calculated by the use of SAS software (Statistical Software System; SAS Institute, Inc. Cary, N.C.).


In Table 1 below, the following ranges apply:

IC₅₀: A = ≥1µM; B = 1µM-500nM; and C<500nM.













EC₅₀: A = ≥1µM; and B = <1µM














76

















TABLE 1



Cpd. #	A	R ²	R ³	Z	IC ₅₀	EC ₅₀	m/z (M+H) ⁺
101	N-Me	phenyl	cyclohexyl	OH	A	--	334.1
102	NH		cyclohexyl	OH	A	A	310.0
103	NH		cyclohexyl	OH	A	--	308.0
104	NH		cyclohexyl	OH	A	--	324.0 (M-H)
105	NH	Br	cyclohexyl	OH	A	--	319.9
106	N-Me		cyclohexyl	OH	B	A	335.2
107	N-Me		cyclohexyl	OH	B	A	324.1
108	N-Me		cyclohexyl	OH	B	B	349.1
109	N-Me		cyclohexyl	OH	C	A	336.1
110	NH		cyclopentyl	OH	C	--	296.0
111	N-Me		cyclopentyl	OH	C	A	310.0
112	N-Me		cyclohexyl	OH	C	A	350.1
113	N-Me		cyclopentyl	OH	C	--	336.1

Cpd. #	A	R ²	R ³	Z	IC ₅₀	EC ₅₀	m/z (M+H) ⁺
114			cyclohexyl	OMe	A	A	382
115	N-Me		cyclopentyl	OH	B	A	321
116			cyclohexyl	OH	C	--	368.1
117	N-Me		cyclopentyl	OH	C	A	327.1
118	N-Me		cyclopentyl	OH	C	A	384.1
119	N-Me		cyclopentyl	OH	B	A	356.2
120	N-Me		cyclopentyl	OH	A	--	370.2
121	N-Me		cyclopentyl	OH	B	A	341.1
122	N-Me		cyclopentyl	OH	A	--	384.2
123	N-Me		cyclopentyl	OH	C	A	400.2

Cpd. #	A	R ²	R ³	Z	IC ₅₀	EC ₅₀	m/z (M+H) ⁺
124	N-Me		cyclopentyl	OH	A	--	384.1
125	N-Me		cyclopentyl	OH	A	--	440.2
126	N-Me		cyclopentyl	OH	A	--	426.2
127	N-Me		cyclopentyl	OH	C	A	413.2
128	N-Me		cyclopentyl	OH	C	A	311.1
129	N-Me		cyclopentyl	OH	B	A	327.1
130	N-Me		cyclopentyl	OH	A	--	335.2
131	N-Me		cyclopentyl	OH	B	A	335.2
132	N-Me		cyclopentyl	OH	C	A	326.1
133	N-Me		cyclopentyl	OH	B	A	335.2
134	N-Me		cyclopentyl	OH	B	--	335.2
135	N-Me		cyclopentyl	OH	C	A	322.2
136	N-Me		cyclopentyl	OH	B	--	399.1

Cpd. #	A	R ²	R ³	Z	IC ₅₀	EC ₅₀	m/z (M+H) ⁺
137	N-Me		cyclopentyl	OH	B	--	366.1
138	S		cyclopentyl	OH	A	A	324.1
139	N-Me		cyclohexyl	OH	C	--	356.1
140	S		cyclopentyl	OH	A	--	331.1
141	O		cyclopentyl	OH	A	A	308.2
142	NH		cyclohexyl	OH	A	--	321.1
143			cyclohexyl	OH	B	--	379.2
144	N-Me		cyclopentyl	OH	A	--	355.0
145	NH		cyclopentyl	OH	A	A	307.1
146			cyclohexyl	OH	A	--	471.1
147	N-Me		cyclopentyl	OH	A	--	351.1
148	N-Me		cyclopentyl	OH	B	--	339.1
149			cyclohexyl	OH	B	--	457.2

80





Cpd. #	A	R ²	R ³	Z	IC ₅₀	EC ₅₀	m/z (M+H) ⁺
150	N-Me			OH	--	--	319.0

TABLE 2




Cpd. #	A	R ²	R ³	Z	IC ₅₀	EC ₅₀	m/z (M+H) ⁺
201	N-Me	phenyl	cyclohexyl	OH	A	--	335.3
202	N-Me		cyclohexyl	OH	A	--	325.2

81

CLAIMS

WHAT IS CLAIMED IS:

1. An isomer, enantiomer, diastereoisomer, or tautomer of a compound, represented by formula I:



wherein:

A is O, S, NR¹, or CR¹, wherein **R**¹ is selected from the group consisting of: H, (C₁₋₆)alkyl optionally substituted with:

- halogen, OR¹¹, SR¹¹ or N(R¹²)₂, wherein **R**¹¹ and each **R**¹² is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-aryl or (C₁₋₆)alkyl-**Het**, said aryl or **Het** optionally substituted with **R**¹⁰; or
- both **R**¹² are covalently bonded together and to the nitrogen to which they are both attached to form a 5, 6 or 7-membered saturated heterocycle;

----- represents either a single or a double bond;

R² is selected from: H, halogen, **R**²¹, OR²¹, SR²¹, COOR²¹, SO₂N(**R**²²)₂, N(**R**²²)₂, CON(**R**²²)₂, NR²²C(O)**R**²² or NR²²C(O)NR²² wherein **R**²¹ and each **R**²² is independently H, (C₁₋₆)alkyl, haloalkyl, (C₂₋₆)alkenyl, (C₃₋₇)cycloalkyl, (C₂₋₆)alkynyl, (C₅₋₇)cycloalkenyl, 6 or 10-membered aryl or **Het**, said **R**²¹ and **R**²² being optionally substituted with **R**²⁰, or both **R**²² are bonded together to form a 5, 6 or 7-membered saturated heterocycle with the nitrogen to which they are attached;

wherein **R**¹⁰ and **R**²⁰ is each:

- 1 to 4 substituents selected from: halogen, OPO₃H, NO₂, cyano, azido, C(=NH)NH₂, C(=NH)NH(C₁₋₆)alkyl or C(=NH)NHCO(C₁₋₆)alkyl; or
- 1 to 4 substituents selected from:
 - a) (C₁₋₆) alkyl or haloalkyl, (C₃₋₇)cycloalkyl, C₃₋₇ spirocycloalkyl optionally containing 1 or 2 heteroatom, (C₂₋₆)alkenyl, (C₃₋₆)cycloalkenyl, (C₂₋₈)alkynyl, (C₁₋₆) alkyl-(C₃₋₇)cycloalkyl, all of which optionally substituted with **R**¹⁵⁰;
 - b) OR¹⁰⁴ wherein **R**¹⁰⁴ is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋

- 7)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said alkyl, cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being optionally substituted with R¹⁵⁰;
- c) OCOR¹⁰⁵ wherein R¹⁰⁵ is (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said alkyl, cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being optionally substituted with R¹⁵⁰;
- d) SR¹⁰⁸, SO₂N(R¹⁰⁸)₂ or SO₂N(R¹⁰⁸)C(O)R¹⁰⁸ wherein each R¹⁰⁸ is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** or both R¹⁰⁸ are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** or heterocycle being optionally substituted with R¹⁵⁰;
- e) NR¹¹¹R¹¹² wherein R¹¹¹ is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, and R¹¹² is H, CN, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl, (C₁₋₆alkyl)**Het**, COOR¹¹⁵ or SO₂R¹¹⁵ wherein R¹¹⁵ is (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, or both R¹¹¹ and R¹¹² are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, or heterocycle being optionally substituted with R¹⁵⁰;
- f) NR¹¹⁶COR¹¹⁷ wherein R¹¹⁶ and R¹¹⁷ is each H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being optionally substituted with R¹⁵⁰;
- g) NR¹¹⁸CONR¹¹⁹R¹²⁰, wherein R¹¹⁸, R¹¹⁹ and R¹²⁰ is each H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, or R¹¹⁸ is covalently bonded to R¹¹⁹ and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle; or R¹¹⁹ and R¹²⁰ are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle; said alkyl, cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** or heterocycle being optionally substituted with R¹⁵⁰;

- h)** $\text{NR}^{121}\text{COCOR}^{122}$ wherein R^{121} and R^{122} is each H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, a 6- or 10-membered aryl, **Het**, $(\text{C}_{1-6})\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6})\text{alkyl})\text{Het}$, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, **Het**, $(\text{C}_{1-6})\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6})\text{alkyl})\text{Het}$ being optionally substituted with R^{150} ; or R^{122} is OR^{123} or $\text{N}(\text{R}^{124})_2$ wherein R^{123} and each R^{124} is independently H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, or $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6})\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6})\text{alkyl})\text{Het}$, or R^{124} is OH or $\text{O}(\text{C}_{1-6})\text{alkyl}$ or both R^{124} are covalently bonded together to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, **Het**, $(\text{C}_{1-6})\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6})\text{alkyl})\text{Het}$ and heterocycle being optionally substituted with R^{150} ;
- i)** COR^{127} wherein R^{127} is H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$ or $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6})\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6})\text{alkyl})\text{Het}$, said alkyl, cycloalkyl, aryl, **Het**, $(\text{C}_{1-6})\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6})\text{alkyl})\text{Het}$ being optionally substituted with R^{150} ;
- j)** COOR^{128} wherein R^{128} is H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, or $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6})\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6})\text{alkyl})\text{Het}$, said $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, or $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6})\text{alkyl})\text{aryl}$ and $(\text{C}_{1-6})\text{alkyl})\text{Het}$ being optionally substituted with R^{150} ;
- k)** $\text{CONR}^{129}\text{R}^{130}$ wherein R^{129} and R^{130} are independently H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6})\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6})\text{alkyl})\text{Het}$, or both R^{129} and R^{130} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, **Het**, $(\text{C}_{1-6})\text{alkyl})\text{aryl}$, $(\text{C}_{1-6})\text{alkyl})\text{Het}$ and heterocycle being optionally substituted with R^{150} ;
- l)** aryl, **Het**, $(\text{C}_{1-6})\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6})\text{alkyl})\text{Het}$, all of which being optionally substituted with R^{150} ; and

wherein R^{150} is defined as:

- 1 to 3 substituents selected from: halogen, OPO_3H , NO_2 , cyano, azido, $\text{C}(=\text{NH})\text{NH}_2$, $\text{C}(=\text{NH})\text{NH}(\text{C}_{1-6})\text{alkyl}$ or $\text{C}(=\text{NH})\text{NHCO}(\text{C}_{1-6})\text{alkyl}$;
or

- 1 to 3 substituents selected from:

a) $(\text{C}_{1-6})\text{alkyl}$ or haloalkyl, $(\text{C}_{3-7})\text{cycloalkyl}$, C_{3-7} spirocycloalkyl optionally containing 1 or 2 heteroatom, $(\text{C}_{2-6})\text{alkenyl}$, $(\text{C}_{2-8})\text{alkynyl}$, $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, all of which optionally substituted with R^{160} ;

b) OR^{104} wherein R^{104} is H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, or $(\text{C}_{1-6})\text{alkyl}-$

(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said alkyl, cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being optionally substituted with R¹⁶⁰;

c) OCOR¹⁰⁵ wherein R¹⁰⁵ is (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said alkyl, cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being optionally substituted with R¹⁶⁰;

d) SR¹⁰⁸, SO₂N(R¹⁰⁸)₂ or SO₂N(R¹⁰⁸)C(O)R¹⁰⁸ wherein each R¹⁰⁸ is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** or both R¹⁰⁸ are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** or heterocycle being optionally substituted with R¹⁶⁰;

e) NR¹¹¹R¹¹² wherein R¹¹¹ is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, and R¹¹² is H, CN, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl, (C₁₋₆alkyl)**Het**, COOR¹¹⁵ or SO₂R¹¹⁵ wherein R¹¹⁵ is (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, or both R¹¹¹ and R¹¹² are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, or heterocycle being optionally substituted with R¹⁶⁰;

f) NR¹¹⁶COR¹¹⁷ wherein R¹¹⁶ and R¹¹⁷ is each H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆)alkyl)**Het**, said (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being optionally substituted with R¹⁶⁰;

g) NR¹¹⁸CONR¹¹⁹R¹²⁰, wherein R¹¹⁸, R¹¹⁹ and R¹²⁰ is each H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆)alkyl)aryl or (C₁₋₆alkyl)**Het**, or R¹¹⁸ is covalently bonded to R¹¹⁹ and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, or R¹¹⁹ and R¹²⁰ are covalently bonded

together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** or heterocycle being optionally substituted with **R**¹⁶⁰;

h) **NR**¹²¹**COCOR**¹²² wherein **R**¹²¹ and **R**¹²² is each H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, a 6- or 10-membered aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being optionally substituted with **R**¹⁶⁰, or **R**¹²² is **OR**¹²³ or **N(R**¹²⁴**)**₂ wherein **R**¹²³ and each **R**¹²⁴ is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, or **R**¹²⁴ is OH or O(C₁₋₆alkyl) or both **R**¹²⁴ are covalently bonded together to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** and heterocycle being optionally substituted with **R**¹⁶⁰;

i) **COR**¹²⁷ wherein **R**¹²⁷ is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said alkyl, cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het** being optionally substituted with **R**¹⁶⁰;

j) tetrazole, **COOR**¹²⁸ wherein **R**¹²⁸ is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, said (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl and (C₁₋₆alkyl)**Het** being optionally substituted with **R**¹⁶⁰; and

k) **CONR**¹²⁹**R**¹³⁰ wherein **R**¹²⁹ and **R**¹³⁰ are independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl or (C₁₋₆alkyl)**Het**, or both **R**¹²⁹ and **R**¹³⁰ are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, **Het**, (C₁₋₆alkyl)aryl, (C₁₋₆alkyl)**Het** and heterocycle being optionally substituted with **R**¹⁶⁰;

wherein **R**¹⁶⁰ is defined as 1 or 2 substituents selected from: tetrazole, halogen, CN, C₁₋₆alkyl, haloalkyl, **COOR**¹⁶¹, **SO**₃H, **SR**¹⁶¹, **SO**₂**R**¹⁶¹, **OR**¹⁶¹, **N(R**¹⁶²**)**₂, **SO**₂**N(R**¹⁶²**)**₂, **NR**¹⁶²**COR**¹⁶² or

CON(R¹⁶²)₂, wherein R¹⁶¹ and each R¹⁶² is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl; or both R¹⁶² are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle,

B is NR³ or CR³, with the proviso that one of **A** or **B** is either CR¹ or CR³, wherein R³ is selected from (C₁₋₆)alkyl, haloalkyl, (C₃₋₇)cycloalkyl, (C₅₋₇)cycloalkenyl, (C₆₋₁₀)bicycloalkyl, (C₆₋₁₀)bicycloalkenyl, 6- or 10-membered aryl, **Het**, (C₁₋₆)alkyl-aryl or (C₁₋₆)alkyl-**Het**,

said alkyl, cycloalkyl, bicycloalkyl, aryl, **Het**, alkyl-aryl and alkyl-**Het** being optionally substituted with from 1 to 4 substituents selected from: halogen, or

a) (C₁₋₆)alkyl optionally substituted with:

- OR³¹ or SR³¹ wherein R³¹ is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆)alkyl-aryl or (C₁₋₆)alkyl-**Het**; or

- N(R³²)₂ wherein each R³² is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆)alkyl-aryl or (C₁₋₆)alkyl-**Het**; or both R³² are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle;

b) OR³³ wherein R³³ is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆)alkyl-aryl or (C₁₋₆)alkyl-**Het**;

c) SR³⁴ wherein R³⁴ is H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆)alkyl-aryl or (C₁₋₆)alkyl-**Het**;

and
d) N(R³⁵)₂ wherein each R³⁵ is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, aryl, **Het**, (C₁₋₆)alkyl-aryl or (C₁₋₆)alkyl-**Het**; or both R³⁵ are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle;

K is N or CR⁴, wherein R⁴ is H, halogen, (C₁₋₆)alkyl, haloalkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl; or R⁴ is OR⁴¹ or SR⁴¹, COR⁴¹ or NR⁴¹COR⁴¹ wherein each R⁴¹

87

is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl; or R⁴ is NR⁴²R⁴³ wherein R⁴² and R⁴³ are each independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl, or both R⁴² and R⁴³ are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle;

L is N or CR⁵, wherein R⁵ has the same definition as R⁴ defined above;


M is N or CR⁷, wherein R⁷ has the same definition as R⁴ defined above;

Y¹ is O or S;

Z is OR⁶, wherein R⁶ is H, (C₁₋₆)alkyl being optionally substituted with: halo, hydroxy, carboxy, amino, C₁₋₆ alkoxy, C₁₋₆alkoxycarbonyl, and C₁₋₆ alkylamino; or R⁶ is C₁₋₆ alkylaryl optionally substituted with: halogen, cyano, nitro, C₁₋₆ alkyl, C₁₋₆haloalkyl, C₁₋₆alkanoyl, -(CH₂)₁₋₆-COOR⁷, -(CH₂)₁₋₆-CONR⁷R⁸, -(CH₂)₁₋₆-NR⁷R⁸, -(CH₂)₁₋₆-NR⁷COR⁸, -(CH₂)₁₋₆-NHSO₂R⁷, -(CH₂)₁₋₆-OR⁷, -(CH₂)₁₋₆-SR⁷, -(CH₂)₁₋₆-SO₂R⁷, and -(CH₂)₁₋₆-SO₂NR⁷R⁸, wherein each R⁷ and each R⁸ is H or C₁₋₆ alkyl,

or Z is NR⁹R¹⁰ wherein each of R⁹ and R¹⁰ is selected from: H, C₁₋₆alkoxy, or C₁₋₆alkyl optionally substituted with halo, hydroxy, carboxy, amino, C₁₋₆ alkoxy, C₁₋₆alkoxycarbonyl, and C₁₋₆ alkylamino;

or a salt thereof;


with the proviso that when A is CR¹, R¹ is Me, R² is pyridine or , B is NR³, R³ is Me, K, L, M is CH, Y¹ is O, and Z is OR⁶, then R⁶ is not H;

and with the proviso that when A is NR¹, R¹ is H, R² is phenyl, B is CR³, R³ is phenyl, K, L, M is CH, Y¹ is O, and Z is OR⁶, then R⁶ is not H;

and with the proviso that when A is S, R² is bromine, B is CR³, R³ is Me, K is CH, L is CH, M is CR⁷, R⁷ is H or Me, Y¹ is O, and Z is OR⁶, then R⁶ is not H;

and with the proviso that when **A** is O, **R**² is H, **B** is CR³, **R**³ is phenyl, **K**, **L**, **M** is CH, **Y**¹ is O, and **Z** is OR⁶, then **R**⁶ is not H;


and with the proviso that when **A** is CR¹, **R**¹ is Me, **R**² is pyridine, **B** is NR³, **R**³ is Me, **K**, **L**, **M** is CH, **Y**¹ is O, and **Z** is OR⁶, then **R**⁶ is not Me;

and with the further proviso that when **A** is CR¹, **R**¹ is Me, **R**² is , **B** is NR³, **R**³ is Me, **K**, **L**, **M** is CH, **Y**¹ is O, and **Z** is OR⁶, then **R**⁶ is not Et;

and with the further proviso that when **A** is CR¹, **R**¹ is CH, **R**² is Me, **B** is NR³, **R**³ is Me, **K**, **L**, **M** is CH, **Y**¹ is O, and **Z** is OR⁶, then **R**⁶ is not Et;


and with the further proviso that when **A** is CR¹, **R**¹ is Et, **R**² is Me, **B** is NR³, **R**³ is Me, **K**, **L**, **M** is CH, **Y**¹ is O, and **Z** is OR⁶, then **R**⁶ is not CH₂CH₂N(Me)₂;

and with the further proviso that when **A** is CH, **R**² is Me, **B** is NR³, **R**³ is



, **K** is N, **L** is CR⁵, **R**⁵ is Me, **M** is CR⁷, **R**⁷ is OH, **Y**¹ is O, and **Z** is OR⁶ then **R**⁶ is not Et;

and with the further proviso that when **A** is NR¹, **R**¹ is Me, **R**² is Br, **B** is CR³, **R**³ is




, **K** is N, **L** is CR⁵, **R**⁵ is Me, **M** is CR⁷, **R**⁷ is Br, **Y**¹ is O, and **Z** is OR⁶, then **R**⁶ is not Me;

and with the further proviso that when **A** is NR¹, **R**¹ is H, **R**² is Cl, **B** is CR³, **R**³ is Et,


89

K is CH, L is CH, M is CH, Y¹ is O, Z is OR⁶, then R⁶ is not Me;

and with the further proviso that when A is NR¹, R¹ is H, R² is phenyl, B is CR³, R³ is phenyl, K is CH, L is CH, M is CR⁷, R⁷ is Me, Y¹ is O, Z is OR⁶, then R⁶ is not Et;




and with the further proviso that when A is NR¹, R¹ is H, R² is



is OMe, K is CH, L is N, M is CH, Y¹ is O, and Z is OR⁶, then R⁶ is not Et;


with the further proviso that when A is S, R² is Br, B is CR³, R³ is Me, K is CH, L is CH, M is CH, Y¹ is O, and Z is OR⁶, then R⁶ is not Me;

and with the further proviso that, when A is NR¹, R¹ is H, R² is:




, B is NR³, R³ is cyclohexyl, K, L, M is CH, Y¹ is O, Z is OR⁶, then R⁶ is not H.

2. The compound according to claim 1, having the following formula (II):




wherein, preferably, A is O, S, or NR¹.

3. The compound according to claim 2, wherein **A** is NR^1 .
4. The compound according to claim 1, wherein having the following formula (III):



wherein, preferably, **B** is NR^3 .

5. The compound according to claim 1, wherein **M**, **K** and **L** is CH or N.
6. The compound according to claim 5, wherein **M**, **K** and **L** is CH.
7. The compound according to claim 1, having the following formulae:



wherein R^1 , R^2 , R^3 and **Z** are as defined in claim 1.

8. The compound according to claim 1, wherein R^1 is selected from the group consisting of: H or (C_{1-6}) alkyl.
9. The compound according to claim 8, wherein R^1 is H, CH_3 , isopropyl, or isobutyl.
10. The compound according to claim 9, wherein R^1 is H or CH_3 .
11. The compound according to claim 10, wherein R^1 is CH_3 .
12. The compound according to claim 1, wherein R^2 is selected from: H, halogen, (C_{2-6}) alkenyl, (C_{5-7}) cycloalkenyl, 6 or 10-membered aryl or **Het**; wherein (C_{2-6}) alkenyl, (C_{5-7}) cycloalkenyl, aryl or **Het** is optionally substituted with R^{20} , wherein R^{20} is defined as:
 - 1 to 4 substituents selected from: halogen, NO_2 , cyano, azido, $C(=NH)NH_2$, $C(=NH)NH(C_{1-6})$ alkyl or $C(=NH)NHCO(C_{1-6})$ alkyl; or
 - 1 to 4 substituents selected from:
 - a) (C_{1-6}) alkyl or haloalkyl, (C_{3-7}) cycloalkyl, (C_{2-6}) alkenyl, (C_{2-8}) alkynyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, all of which optionally substituted with R^{150} ;
 - b) OR^{104} wherein R^{104} is H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, or (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het**, said alkyl, cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het** being optionally substituted with R^{150} ;
 - c) $OCOR^{105}$ wherein R^{105} is (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl, (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het**, said alkyl, cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het** being optionally substituted with R^{150} ;
 - d) SR^{108} , $SO_2N(R^{108})_2$ or $SO_2N(R^{108})C(O)R^{108}$ wherein each R^{108} is independently H, (C_{1-6}) alkyl, (C_{3-7}) cycloalkyl or (C_{1-6}) alkyl- (C_{3-7}) cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het** or both R^{108} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, aryl, **Het**, (C_{1-6}) alkyl)aryl or (C_{1-6}) alkyl)**Het** or heterocycle being optionally substituted with R^{150} ;

- e) $\text{NR}^{111}\text{R}^{112}$ wherein R^{111} is H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$ or $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6}\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6}\text{alkyl})\text{Het}$, and R^{112} is H, CN, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$ or $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6}\text{alkyl})\text{aryl}$, $(\text{C}_{1-6}\text{alkyl})\text{Het}$, COOR^{115} or $\text{SO}_2\text{R}^{115}$ wherein R^{115} is $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, or $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6}\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6}\text{alkyl})\text{Het}$, or both R^{111} and R^{112} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, aryl, **Het**, $(\text{C}_{1-6}\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6}\text{alkyl})\text{Het}$, or heterocycle being optionally substituted with R^{150} ;
- f) $\text{NR}^{116}\text{COR}^{117}$ wherein R^{116} and R^{117} is each H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6}\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6}\text{alkyl})\text{Het}$, said $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6}\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6}\text{alkyl})\text{Het}$ being optionally substituted with R^{150} ;
- g) $\text{NR}^{118}\text{CONR}^{119}\text{R}^{120}$, wherein R^{118} , R^{119} and R^{120} is each H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6}\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6}\text{alkyl})\text{Het}$, or R^{118} is covalently bonded to R^{119} and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle; or R^{119} and R^{120} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle; said alkyl, cycloalkyl, $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6}\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6}\text{alkyl})\text{Het}$ or heterocycle being optionally substituted with R^{150} ;
- h) $\text{NR}^{121}\text{COCOR}^{122}$ wherein R^{121} and R^{122} is each H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, a 6- or 10-membered aryl, **Het**, $(\text{C}_{1-6}\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6}\text{alkyl})\text{Het}$, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, **Het**, $(\text{C}_{1-6}\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6}\text{alkyl})\text{Het}$ being optionally substituted with R^{150} ; or R^{122} is OR^{123} or $\text{N}(\text{R}^{124})_2$ wherein R^{123} and each R^{124} is independently H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, or $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6}\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6}\text{alkyl})\text{Het}$, or R^{124} is OH or $\text{O}(\text{C}_{1-6}\text{alkyl})$ or both R^{124} are covalently bonded together to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, **Het**, $(\text{C}_{1-6}\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6}\text{alkyl})\text{Het}$ and heterocycle being optionally substituted with R^{150} ;
- i) COR^{127} wherein R^{127} is H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$ or $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6}\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6}\text{alkyl})\text{Het}$, said alkyl, cycloalkyl, aryl, **Het**, $(\text{C}_{1-6}\text{alkyl})\text{aryl}$ or $(\text{C}_{1-6}\text{alkyl})\text{Het}$ being optionally substituted with R^{150} ;

j) COOR^{128} wherein R^{128} is H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, or $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6})\text{alkyl}$ aryl or $(\text{C}_{1-6})\text{alkyl}$ **Het**, said $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, or $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6})\text{alkyl}$ aryl and $(\text{C}_{1-6})\text{alkyl}$ **Het** being optionally substituted with R^{150} ;

k) $\text{CONR}^{129}\text{R}^{130}$ wherein R^{129} and R^{130} are independently H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6})\text{alkyl}$ aryl or $(\text{C}_{1-6})\text{alkyl}$ **Het**, or both R^{129} and R^{130} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, alkyl-cycloalkyl, aryl, **Het**, $(\text{C}_{1-6})\text{alkyl}$ aryl, $(\text{C}_{1-6})\text{alkyl}$ **Het** and heterocycle being optionally substituted with R^{150} ;

l) aryl, **Het**, $(\text{C}_{1-6})\text{alkyl}$ aryl or $(\text{C}_{1-6})\text{alkyl}$ **Het**, all of which being optionally substituted with R^{150} , wherein R^{150} is preferably:

- 1 to 3 substituents selected from: halogen, NO_2 , cyano or azido; or

- 1 to 3 substituents selected from:

a) $(\text{C}_{1-6})\text{alkyl}$ or haloalkyl, $(\text{C}_{3-7})\text{cycloalkyl}$, $(\text{C}_{2-6})\text{alkenyl}$, $(\text{C}_{2-8})\text{alkynyl}$, $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, all of which optionally substituted with R^{160} ;

b) OR^{104} wherein R^{104} is H, $(\text{C}_{1-6})\text{alkyl}$ or $(\text{C}_{3-7})\text{cycloalkyl}$, said alkyl or cycloalkyl optionally substituted with R^{160} ;

d) SR^{108} , $\text{SO}_2\text{N}(\text{R}^{108})_2$ or $\text{SO}_2\text{N}(\text{R}^{108})\text{C}(\text{O})\text{R}^{108}$ wherein each R^{108} is independently H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$ or $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, or both R^{108} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, aryl, **Het** and heterocycle being optionally substituted with R^{160} ;

e) $\text{NR}^{111}\text{R}^{112}$ wherein R^{111} is H, $(\text{C}_{1-6})\text{alkyl}$, or $(\text{C}_{3-7})\text{cycloalkyl}$, and R^{112} is H, $(\text{C}_{1-6})\text{alkyl}$ or $(\text{C}_{3-7})\text{cycloalkyl}$, COOR^{115} or $\text{SO}_2\text{R}^{115}$ wherein R^{115} is $(\text{C}_{1-6})\text{alkyl}$ or $(\text{C}_{3-7})\text{cycloalkyl}$, or both R^{111} and R^{112} are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl and heterocycle being optionally substituted with R^{160} ;

f) $\text{NR}^{116}\text{COR}^{117}$ wherein R^{116} and R^{117} is each H, $(\text{C}_{1-6})\text{alkyl}$ or $(\text{C}_{3-7})\text{cycloalkyl}$ said $(\text{C}_{1-6})\text{alkyl}$ and $(\text{C}_{3-7})\text{cycloalkyl}$ being optionally substituted with R^{160} ;

g) $\text{NR}^{118}\text{CONR}^{119}\text{R}^{120}$, wherein R^{118} , R^{119} and R^{120} is each H, $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{3-7})\text{cycloalkyl}$, $(\text{C}_{1-6})\text{alkyl}-(\text{C}_{3-7})\text{cycloalkyl}$, aryl, **Het**, $(\text{C}_{1-6})\text{alkyl}$ aryl or $(\text{C}_{1-6})\text{alkyl}$ **Het**, all of which being optionally substituted with R^{150} , wherein R^{150} is preferably:

₆)alkyl or (C₃₋₇)cycloalkyl, or R¹¹⁸ is covalently bonded to R¹¹⁹ and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, or R¹¹⁹ and R¹²⁰ are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl, and heterocycle being optionally substituted with R¹⁶⁰;

h) NR¹²¹COCOR¹²² wherein R¹²¹ is H, (C₁₋₆)alkyl or (C₃₋₇)cycloalkyl, said alkyl and cycloalkyl being optionally substituted with R¹⁶⁰, or R¹²² is OR¹²³ or N(R¹²⁴)₂ wherein R¹²³ and each R¹²⁴ is independently H, (C₁₋₆)alkyl or (C₃₋₇)cycloalkyl, or both R¹²⁴ are covalently bonded together to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl and heterocycle being optionally substituted with R¹⁶⁰;

i) COR¹²⁷ wherein R¹²⁷ is H, (C₁₋₆)alkyl or (C₃₋₇)cycloalkyl, said alkyl and cycloalkyl being optionally substituted with R¹⁶⁰;

j) COOR¹²⁸ wherein R¹²⁸ is H, (C₁₋₆)alkyl or (C₃₋₇)cycloalkyl, said (C₁₋₆)alkyl and (C₃₋₇)cycloalkyl being optionally substituted with R¹⁶⁰; and

k) CONR¹²⁹R¹³⁰ wherein R¹²⁹ and R¹³⁰ are independently H, (C₁₋₆)alkyl or (C₃₋₇)cycloalkyl, or both R¹²⁹ and R¹³⁰ are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle, said alkyl, cycloalkyl and heterocycle being optionally substituted with R¹⁶⁰;

wherein R¹⁶⁰ is defined as 1 or 2 substituents selected from: halogen, CN, C₁₋₆alkyl, haloalkyl, COOR¹⁶¹, OR¹⁶¹, N(R¹⁶²)₂, SO₂N(R¹⁶²)₂, NR¹⁶²COR¹⁶² or CON(R¹⁶²)₂, wherein R¹⁶¹ and each R¹⁶² is independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl or (C₁₋₆)alkyl-(C₃₋₇)cycloalkyl; or both R¹⁶² are covalently bonded together and to the nitrogen to which they are attached to form a 5, 6 or 7-membered saturated heterocycle.

13. The compound according to claim 12, wherein R² is selected from: aryl or Het, each optionally monosubstituted or disubstituted with substituents selected from the group consisting of: halogen, haloalkyl, N₃, or


a) (C₁₋₆)alkyl optionally substituted with OH, O(C₁₋₆)alkyl or SO₂(C₁₋₆

- alkyl);
- b) (C₁₋₆)alkoxy;
 - e) $\text{NR}^{111}\text{R}^{112}$ wherein both R^{111} and R^{112} are independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or R^{112} is 6- or 10-membered aryl, **Het**, (C₁₋₆)alkyl-aryl or (C₁₋₆)alkyl-**Het**; or both R^{111} and R^{112} are covalently bonded together and to the nitrogen to which they are attached to form a nitrogen-containing heterocycle, each of said alkyl, cycloalkyl, aryl, **Het**, alkyl-aryl or alkyl-**Het**; being optionally substituted with halogen or:
 - OR^{161} or $\text{N}(\text{R}^{162})_2$, wherein R^{161} and each R^{162} is independently H, (C₁₋₆)alkyl, or both R^{162} are covalently bonded together and to the nitrogen to which they are attached to form a nitrogen-containing heterocycle;
 - f) NHCOR^{117} wherein R^{117} is (C₁₋₆)alkyl, O(C₁₋₆)alkyl or O(C₃₋₇)cycloalkyl;
 - i) CO-aryl; and
 - k) CONH_2 , $\text{CONH}(\text{C}_{1-6}\text{alkyl})$, $\text{CON}(\text{C}_{1-6}\text{alkyl})_2$, CONH-aryl , or $\text{CONHC}_{1-6}\text{alkyl-aryl}$.


14. The compound according to claim 13, wherein R^2 is aryl or **Het**, each optionally monosubstituted or disubstituted with substituents selected from the group consisting of: halogen, haloalkyl, or

- a) (C₁₋₆)alkyl optionally substituted with OH, O(C₁₋₆)alkyl or $\text{SO}_2(\text{C}_{1-6}\text{alkyl})$;
- b) (C₁₋₆)alkoxy; and
- e) $\text{NR}^{111}\text{R}^{112}$ wherein both R^{111} and R^{112} are independently H, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, or R^{112} is 6- or 10-membered aryl, **Het**, (C₁₋₆)alkyl-aryl or (C₁₋₆)alkyl-**Het**; or both R^{111} and R^{112} are covalently bonded together and to the nitrogen to which they are attached to form a nitrogen-containing heterocycle, each of said alkyl, cycloalkyl, aryl, **Het**, alkyl-aryl or alkyl-**Het**; or being optionally substituted with halogen or:
 - OR^{161} or $\text{N}(\text{R}^{162})_2$, wherein R^{161} and each R^{162} is independently H, (C₁₋₆)alkyl, or both R^{162} are covalently bonded together and to the nitrogen to which they are attached to form a nitrogen-containing heterocycle.


15. The compound according to claim 14, wherein R^2 is phenyl or a heterocycle selected from:




16. The compound according to claim 15, wherein R^2 is selected from:
H, Br, $CH=CH_2$,




97




98




17. The compound according to claim 16, wherein R^2 is selected from:





18. The compound according to claim 17, wherein R^2 is selected from:



19. The compound according to claim 1, wherein R^3 is selected from (C_{3-7}) cycloalkyl, (C_{3-7}) cycloalkenyl, (C_{6-10}) bicycloalkyl, (C_{6-10}) bicycloalkenyl, 6- or 10-membered aryl, or **Het**.

20. The compound according to claim 19, wherein R^3 is (C_{3-7}) cycloalkyl.

21. The compound according to claim 20, wherein R^3 is cyclopentyl, or cyclohexyl.

22. The compound according to claim 1, wherein Y^1 is O.

23. The compound according to claim 1, wherein Z is OR^6 , wherein R^6 is H, (C_{1-6}) alkyl being optionally substituted with: halo, hydroxy, carboxy, amino, C_{1-6} alkoxy, C_{1-6} alkoxycarbonyl, and C_{1-6} alkylamino; or R^6 is C_{1-6} alkylaryl optionally substituted with: halogen, cyano, nitro, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkanoyl, $-(CH_2)_1-$

100


$-COOR^7$, $-(CH_2)_{1-6}-CONR^7R^8$, $-(CH_2)_{1-6}-NR^7R^8$, $-(CH_2)_{1-6}-NR^7COR^8$, $-(CH_2)_{1-6}-NHSO_2R^7$, $-(CH_2)_{1-6}-OR^7$, $-(CH_2)_{1-6}-SR^7$, $-(CH_2)_{1-6}-SO_2R^7$, and $-(CH_2)_{1-6}-SO_2NR^7R^8$, wherein each R^7 and each R^8 is H or C_{1-6} alkyl,

or Z is NR^9R^{10} wherein each of R^9 and R^{10} is selected from: H, C_{1-6} alkoxy, or C_{1-6} alkyl optionally substituted with halo, hydroxy, carboxy, amino, C_{1-6} alkoxy, C_{1-6} alkoxycarbonyl, and C_{1-6} alkylamino.

24. The compound according to claim 23, wherein Z is OH or $O(C_{1-6}alkyl)$ or Z is NR^9R^{10} wherein R^9 is H and R^{10} is H or $C_{1-6}alkyl$.

25. The compound according to claim 24, wherein Z is OH.

26. A compound selected from compounds of formula:













wherein A , R^2 , R^3 and Z are as defined below:

Cpd. #	A	R^2	R^3	Z
101	N-Me	phenyl	cyclohexyl	OH
102	NH		cyclohexyl	OH
103	NH		cyclohexyl	OH
104	NH		cyclohexyl	OH
105	NH	Br	cyclohexyl	OH
106	N-Me		cyclohexyl	OH
107	N-Me		cyclohexyl	OH
















101

Cpd. #	A	R ²	R ³	Z
108	N-Me		cyclohexyl	OH ;
109	N-Me		cyclohexyl	OH ;
110	NH		cyclopentyl	OH ;
111	N-Me		cyclopentyl	OH ;
112	N-Me		cyclohexyl	OH ;
113	N-Me		cyclopentyl	OH ;
114			cyclohexyl	OMe ;
115	N-Me		cyclopentyl	OH ;
116			cyclohexyl	OH ;
117	N-Me		cyclopentyl	OH ;
118	N-Me		cyclopentyl	OH ;
119	N-Me		cyclopentyl	OH ;
120	N-Me		cyclopentyl	OH ;










102

Cpd. #	A	R ²	R ³	Z
121	N-Me		cyclopentyl	OH ;
122	N-Me		cyclopentyl	OH ;
123	N-Me		cyclopentyl	OH ;
124	N-Me		cyclopentyl	OH ;
125	N-Me		cyclopentyl	OH ;
126	N-Me		cyclopentyl	OH ;
127	N-Me		cyclopentyl	OH ;
128	N-Me		cyclopentyl	OH ;
129	N-Me		cyclopentyl	OH ;
130	N-Me		cyclopentyl	OH ;


103

Cpd. #	A	R ²	R ³	Z
131	N-Me		cyclopentyl	OH ;
132	N-Me		cyclopentyl	OH ;
133	N-Me		cyclopentyl	OH ;
134	N-Me		cyclopentyl	OH ;
135	N-Me		cyclopentyl	OH ;
136	N-Me		cyclopentyl	OH ;
137	N-Me		cyclopentyl	OH ;
138	S		cyclopentyl	OH ;
139	N-Me		cyclohexyl	OH ;
140	S		cyclopentyl	OH ;
141	O		cyclopentyl	OH ;
142	NH		cyclohexyl	OH ;
143			cyclohexyl	OH ;
144	N-Me		cyclopentyl	OH ;


104

Cpd. #	A	R ²	R ³	Z
145	NH		cyclopentyl	OH
146			cyclohexyl	OH
147	N-Me		cyclopentyl	OH
148	N-Me		cyclopentyl	OH
149			cyclohexyl	OH
150	N-Me			OH

27. A compound selected from compounds of formula:




wherein A, R², R³ and Z are as defined below:

Cpd. #	A	R ²	R ³	Z
201	N-Me	phenyl	cyclohexyl	OH
202	N-Me		cyclohexyl	OH

105

28. A compound represented by Formula Ia:



wherein:

A is O, S, NR¹, or CR¹;

B is NR³ or CR³;

R¹ is selected from the group consisting of: H, (C₁₋₆)alkyl, benzyl, (C₁₋₆ alkyl)-(C₆₋₁₀aryl), (C₁₋₆ alkyl)-5- or 6-membered heterocycle having 1 to 4 heteroatoms selected from O, N, and S, and 5- or 6-membered heterocycle having 1 to 4 heteroatoms selected from O, N and S,

wherein said benzyl and said heteroatom are optionally substituted with from 1 to 4 substituents selected from the group consisting of: COOH, COO(C₁₋₆ alkyl), halogen, and (C₁₋₆ alkyl);

R² is selected from the group consisting of: H, halogen, (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, phenyl, 5- or 6-membered heterocycle having 1 to 4 heteroatoms selected from O, N, and S, pyridine-N-oxide, and 9- or 10-membered heterobicyclic having 1 to 4 heteroatoms selected from O, N and S,

said phenyl, heterocycle and heterobicyclic being optionally substituted with from 1 to 4 substituents selected from the group consisting of: halogen, C(halogen)₃, (C₁₋₆)alkyl, OH, O(C₁₋₆ alkyl), NH₂, and N(C₁₋₆ alkyl)₂;

R³ is selected from the group consisting of: 5-, 6- or 7-membered heterocycle having 1 to 4 heteroatoms selected from O, N, and S, norbornane, (C₃₋₇)cycloalkyl and (C₃₋₇)cycloalkyl-(C₁₋₆ alkyl);

M is N, CR⁴, or COR⁵, wherein **R⁴** is selected from the group consisting of: H, halogen, and (C₁₋₆ alkyl); and **R⁵** is selected from the group consisting of: H and (C₁₋₆ alkyl);

K and **L** is N or CH;

106

----- represents either a single or a double bond;

Y is O;

Z is OR^6 or NR^6R^{6a} , wherein R^6 is selected from the group consisting of: H, (C_{1-6}) alkyl, wherein said alkyl is optionally substituted with from 1 to 4 substituents selected from: OH, COOH, $COO(C_{1-6})$ alkyl, (C_{1-6}) alkyl, said alkyl being optionally substituted with from 1 to 4 substituents selected from: COOH, $NHCO(C_{1-6})$ alkyl, NH_2 , $NH(C_{1-6})$ alkyl, and $N(C_{1-6})alkyl)_2$;


or a salt thereof.

29. A compound of the formula I according to claim 1, or a pharmaceutically acceptable salt thereof, as an inhibitor of HCV replication.
30. A pharmaceutical composition for the treatment or prevention of HCV infection, comprising an effective amount of a compound of formula I according to claim 1, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.
31. The pharmaceutical composition according to claim 30, further comprising immunomodulatory agent.
32. The pharmaceutical composition according to claim 31, wherein said immunomodulatory agent is selected from: α -, β -, δ -, γ -, and ω -interferons.
33. The pharmaceutical composition according to claim 30, further comprising ribavirin or amantadine.
34. The pharmaceutical composition according to claim 30, further comprising another inhibitor of HCV polymerase.

107


35. The pharmaceutical composition according to claim 34, further comprising an inhibitor of other HCV target, selected from: helicase, polymerase, metalloprotease and IRES.

36. An intermediate of formula (1a) or (1b):



wherein **A**, **B**, **K**, **L**, and **M** are as defined in claim 1, **PG** is H or a carboxy protecting group and **X** is a metal.


37. A process for producing compounds of formula (iii):



wherein **A**, **R²**, **B**, **K**, **L**, **M**, **PG** and **X** are as defined in claim 36, comprising:

a) coupling, in the presence of a metal catalyst, a base and an additive in an appropriate solvent, intermediate (1a) according to claim 36, with **R²-X**.

38. A process for producing compounds of formula (iii),



wherein **A**, **R²**, **B**, **K**, **L**, **M**, and **PG** are as defined in claim 36, comprising:

a) coupling, in the presence of a metal catalyst, a base and an additive in an appropriate solvent, intermediate (1b) according to claim 36, with **R²-X'**, wherein **X'** is as defined in claim 36.

108

39. A process according to claim 37 or 38, wherein said metal catalyst is selected from: Pd, Ni, Ru and Cu.
40. A process according to claim 37 or 38, wherein said additive is selected from: phosphine ligand, Cu salt, Li salt, ammonium salt and CsF.
41. A process according to claim 37 or 38, wherein said metal is selected from: Li, $\text{Sn}(\text{C}_{1-6}\text{alkyl})_3$, $\text{Sn}(\text{aryl})_3$, $\text{B}(\text{OH})_2$, $\text{B}(\text{OC}_{1-6}\text{alkyl})_2$ and metal halide.
42. Use of a compound of formula I according to claim 1, for the manufacture of a medicament for the treatment of HCV infection.

1/3

SEQUENCE LISTING

5 <110> Boehringer Ingelheim (Canada) Ltd.
 <120> Viral Polymerase inhibitors

 10 <130> 13/095
 <140> 60/307,674
 <141> 2001-07-25

 15 <150> 60/338,061
 <151> 2001-12-07

 <160> 4

 20 <170> FastSEQ for Windows Version 4.0
 <210> 1
 <211> 621
 <212> PRT
 <213> HCV NS5B

 25 <400> 1
 Met Ser Tyr Tyr His His His His His Asp Tyr Asp Ile Pro Thr
 1 5 10 15
 Thr Glu Asn Leu Tyr Phe Gln Gly Ala Met Asp Pro Glu Phe Ser Met
 20 25 30
 30 Ser Tyr Thr Trp Thr Gly Ala Leu Ile Thr Pro Cys Ala Ala Glu Glu
 35 40 45
 Ser Gln Leu Pro Ile Asn Ala Leu Ser Asn Ser Leu Val Arg His Arg
 50 55 60
 35 Asn Met Val Tyr Ser Thr Thr Ser Arg Ser Ala Ala Leu Arg Gln Lys
 65 70 75 80
 Lys Val Thr Phe Asp Arg Leu Gln Val Leu Asp Asp His Tyr Arg Asp
 85 90 95
 Val Leu Lys Glu Met Lys Ala Lys Ala Ser Thr Val Lys Ala Lys Leu
 100 105 110
 40 Leu Ser Val Glu Glu Ala Cys Lys Leu Thr Pro Pro His Ser Ala Lys
 115 120 125
 Ser Lys Phe Gly Tyr Gly Ala Lys Asp Val Arg Asn Leu Ser Ser Lys
 130 135 140
 45 Ala Val Asp His Ile Arg Ser Val Trp Lys Asp Leu Leu Glu Asp Thr
 145 150 155 160
 Glu Thr Pro Ile Asp Thr Thr Ile Met Ala Lys Asn Glu Val Phe Cys
 165 170 175
 Val Gln Pro Glu Lys Gly Gly Arg Lys Pro Ala Arg Leu Ile Val Phe
 180 185 190
 50 Pro Asp Leu Gly Val Arg Val Cys Glu Lys Met Ala Leu Tyr Asp Val
 195 200 205
 Val Ser Thr Leu Pro Gln Ala Val Met Gly Ser Ser Tyr Gly Phe Gln
 210 215 220
 55 Tyr Ser Pro Lys Gln Arg Val Glu Phe Leu Val Asn Ala Trp Lys Ser
 225 230 235 240
 Lys Lys Cys Pro Met Gly Phe Ser Tyr Asp Thr Arg Cys Phe Asp Ser
 245 250 255
 Thr Val Thr Glu Ser Asp Ile Arg Val Glu Glu Ser Ile Tyr Gln Cys
 260 265 270
 60 Cys Asp Leu Ala Pro Glu Ala Arg Gln Ala Ile Lys Ser Leu Thr Glu
 275 280 285
 Arg Leu Tyr Ile Gly Gly Pro Leu Thr Asn Ser Lys Gly Gln Asn Cys

SUBSTITUTE SHEET (RULE 26)

2/3

	290		295		300
	Gly Tyr Arg Arg Cys Arg	Ala Ser Gly Val Leu Thr Thr Ser Cys Gly			
	305	310	315	320	
5	Asn Thr Leu Thr Cys Tyr Leu Lys Ala Ser Ala Ala Cys Arg Ala Ala				
	325	330	335		
	Lys Leu Gln Asp Cys Thr Met Leu Val Asn Gly Asp Asp Leu Val Val				
	340	345	350		
	Ile Cys Glu Ser Ala Gly Thr Gln Glu Asp Ala Ala Asn Leu Arg Val				
	355	360	365		
10	Phe Thr Glu Ala Met Thr Arg Tyr Ser Ala Pro Pro Gly Asp Leu Pro				
	370	375	380		
	Gln Pro Glu Tyr Asp Leu Glu Leu Ile Thr Ser Cys Ser Ser Asn Val				
	385	390	395	400	
15	Ser Val Ala His Asp Ala Ser Gly Lys Arg Val Tyr Tyr Leu Thr Arg				
	405	410	415		
	Asp Pro Thr Thr Pro Leu Ala Arg Ala Ala Trp Glu Thr Ala Arg His				
	420	425	430		
	Thr Pro Ile Asn Ser Trp Leu Gly Asn Ile Ile Met Tyr Ala Pro Thr				
	435	440	445		
20	Leu Trp Ala Arg Met Val Leu Met Thr His Phe Phe Ser Ile Leu Leu				
	450	455	460		
	Ala Gln Glu Gln Leu Glu Lys Ala Leu Asp Cys Gln Ile Tyr Gly Ala				
	465	470	475	480	
	Cys Tyr Ser Ile Glu Pro Leu Asp Leu Pro Gln Ile Ile Glu Arg Leu				
25		485	490	495	
	His Gly Leu Ser Ala Phe Ser Leu His Ser Tyr Ser Pro Gly Glu Ile				
	500	505	510		
	Asn Arg Val Ala Ser Cys Leu Arg Lys Leu Gly Val Pro Leu Arg				
	515	520	525		
30	Val Trp Arg His Arg Ala Arg Ser Val Arg Ala Lys Leu Leu Ser Gln				
	530	535	540		
	Gly Gly Arg Ala Ala Thr Cys Gly Lys Tyr Leu Phe Asn Trp Ala Val				
	545	550	555	560	
	Arg Thr Lys Leu Lys Leu Thr Pro Ile Pro Ala Ala Ser Arg Leu Asp				
35		565	570	575	
	Leu Ser Gly Trp Phe Val Ala Gly Tyr Asn Gly Gly Asp Ile Tyr His				
	580	585	590		
	Ser Leu Ser Arg Ala Arg Pro Arg Trp Phe Met Leu Cys Leu Leu Leu				
	595	600	605		
40	Leu Ser Val Gly Val Gly Ile Tyr Leu Leu Pro Asn Arg				
	610	615	620		

45	<210> 2	
	<211> 30	
	<212> DNA	
	<213> Forward Primer	
50	<400> 2	
	acgcagaaag cgtctagcca tggcgtagt	30
	<210> 3	
	<211> 30	
	<212> DNA	
55	<213> Reverse Primer	
	<400> 3	
	tcccggggca ctcgcaagca ccctatcagg	30
60	<210> 4	
	<211> 26	
	<212> DNA	
	<213> PUTR probe	

SUBSTITUTE SHEET (RULE 26)

3/3

<400> 4
tggctcgcgg aaccggtgag tacacc

26

5

SUBSTITUTE SHEET (RULE 26)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
6 February 2003 (06.02.2003)

PCT

(10) International Publication Number
WO 03/010140 A3(51) International Patent Classification⁷: **A61K 31/404**,
31/381, 31/4402, 31/437, C07D 209/08, 405/04, 409/04,
401/04, 403/04, 417/04, 413/04, 471/04[CA/CA]; 2100 Cunard Street, Laval, Québec H7S 2G5
(CA).

(21) International Application Number: PCT/CA02/01127

(74) Agent: **BERNIER, Louise, G.**; Boehringer Ingelheim
(Canada) Ltd., 2100 Cunard Street, Laval, Québec H7S
2G5 (CA).

(22) International Filing Date: 18 July 2002 (18.07.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/307,674 25 July 2001 (25.07.2001) US
60/338,061 7 December 2001 (07.12.2001) US(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).(71) Applicant (*for all designated States except US*):
BOEHRINGER INGELHEIM (CANADA) LTD.
[CA/CA]; 2100 Cunard Street, Laval, Québec H7S 2G5
(CA).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BEAULIEU,**
Pierre, Louis [CA/CA]; 2100 Cunard Street, Laval,
Québec H7S 2G5 (CA). **FAZAL, Gulrez** [CA/CA]; 2100
Cunard Street, Laval, Québec H7S 2G5 (CA). **KUKOLJ,**
George [CA/CA]; 2100 Cunard Street, Laval, Québec H7S
2G5 (CA). **JOLICOEUR, Eric** [CA/CA]; 2100 Cunard
Street, Laval, Québec H7S 2G5 (CA). **GILLARD, James**
[CA/CA]; 2100 Cunard Street, Laval, Québec H7S 2G5
(CA). **POUPART, Marc-André** [CA/CA]; 2100 Cunard
Street, Laval, Québec H7S 2G5 (CA). **RANCOURT, Jean**


Published:

— with international search report

(88) Date of publication of the international search report:
10 July 2003For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 03/010140 A3

(54) Title: HEPATITIS C VIRUS POLYMERASE INHIBITORS WITH HETEROBICYCLIC STRUCTURE

(57) Abstract: An isomer, enantiomer, diastereoisomer, or tautomer of a com-
pound, represented by formula (I) wherein: A is O, S, NR₁, or CR₁, wherein
R₁ is defined herein; represents either a single or a double bond; R₂ is selected
from: H, halogen, R₂₁, OR₂₁, SR₂₁, COOR₂₁, SO₂N(R²²)₂, N(R²²)₂, CON(R²²)₂,
NR²²C(O)R²² or NR²²C(O)NR²² wherein R²¹ and each R²² is defined herein; B
is NR³ or CR³, with the proviso that one of A or B is either CR¹ or CR³, wherein
R³ is defined herein; K is N or CR⁴, wherein R⁴ is defined herein; L is N or CR⁵,
wherein R⁵ has the same definition as R⁴ defined above; M is N or CR⁷, whereinR⁷ has the same definition as R⁴ defined above; Y¹ is O or S; Z is N(R^{6a})R⁶ or OR⁶, wherein R^{6a} is H or alkyl or NR⁶¹R⁶² wherein R⁶¹
and R⁶² are defined herein; a salt or a derivative thereof, as an inhibitor of HCV NS^{5B} polymerase.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 02/01127

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/404 A61K31/381 A61K31/4402 A61K31/437 C07D209/08
 C07D405/04 C07D409/04 C07D401/04 C07D403/04 C07D417/04
 C07D413/04 C07D471/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01 47883 A (JAPAN TOBACCO INC (JP)) 5 July 2001 (2001-07-05) cited in the application abstract page 1 example 503 claims & EP 1 162 196 A1 ---	1-42
X	US 5 059 621 A (C.I.R.D. (FR)) 22 October 1991 (1991-10-22) formulae (I)-(IV), columns 1-3 claims; examples ---	1-28, 36-41
X	US 6 069 156 A (FUJISAWA PHARMACEUTICAL CO LTD (JP)) 30 May 2000 (2000-05-30) formula I, column 1 claims; examples ---	1-28, 36-41
	--- -/-	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

7 March 2003

Date of mailing of the international search report

04/04/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Cortés, J

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 02/01127

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 48697 A (RHONE POULENC RORER LTD (GB)) 24 December 1997 (1997-12-24) claims; examples ---	1-28, 36-41
X	JP 10 067682 A (FUJISAWA PHARMACEUT CO LTD (JP)) 10 March 1998 (1998-03-10) claims; examples ---	1-28, 36-41
X	JP 10 114654 A (FUJISAWA PHARMACEUT CO LTD (JP)) 6 May 1998 (1998-05-06) claims; examples ---	1-28, 36-41
X	JP 09 124632 A (SANKYO CO LTD (JP)) 13 May 1997 (1997-05-13) claims; examples ---	1-28, 36-41
X	WO 01 32653 A (CEPHALON INC (US)) 10 May 2001 (2001-05-10) page 1, line 21 e.g. compounds no. 16, 50, 115, 116, 211, 237-239, 251, 255, 258, ---	1-28, 36-41
X	US 3 565 912 A (UPJOHN CO (US)) 23 February 1971 (1971-02-23) formula IX, examples ---	1-28, 36-41
X	JP 2001 122855 A (JAPAN TOBACCO INC (JP)) 8 May 2001 (2001-05-08) formula (1) e.g. compounds 1-29 to 1-31 ---	1-28, 36-41
X	WO 01 30774 A (AVENTIS PHARMA DEUTSCHLAND GMBH (DE)) 3 May 2001 (2001-05-03) claims; examples ---	1-28, 36-41
X	US 5 912 260 A (JAMES BLACK FOUNDATION LTD. (UK)) 15 June 1999 (1999-06-15) formula (I), examples ---	1-28, 36-41
X	US 4 740 519 A (CENTRE NATIONAL DE RECHERCHES DERMATOLOGIQUES C.I.R.D. (FR)) 26 April 1988 (1988-04-26) formula (I), examples ---	1-28, 36-41
X	EP 0 987 250 A (SHIONOGI & CO LTD (JP)) 22 March 2000 (2000-03-22) e.g. intermediates of formulae (VII) and (VIII) ---	1-28, 36-41
X	WO 99 28297 A (BOEHRINGER INGELHEIM PHARMA (DE)) 10 June 1999 (1999-06-10) claims; examples ---	1-28, 36-41
	--- -/--	

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 02/01127

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 242 167 A (ICI AMERICAS INC (US)) 21 October 1987 (1987-10-21) claims; examples ----	1-28, 36-41
X	WO 01 47922 A (AVENTIS PHARMA LTD (GB)) 5 July 2001 (2001-07-05) claims; examples ----	1-28, 36-41
X	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; HISHMAT, O. H. ET AL: "Synthesis of pharmacologically active indoles" retrieved from STN Database accession no. 132:49850 XP002233311 abstract & BOLLETTINO CHIMICO FARMACEUTICO (1999), 138(6), 259-266, ----	1-28, 36-41
X	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; FUERSTNER, ALOIS ET AL: "Titanium-induced zipper reactions" retrieved from STN Database accession no. 123:83146 XP002233857 abstract & ANGEWANDTE CHEMIE, INTERNATIONAL EDITION IN ENGLISH (1995), 34(6), 678-81, ----	1-28, 36-41
X	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; ROTH, H. J. ET AL: "Synthesis of indole and carbazole derivatives by condensation of alpha.-hydroxyketones and aromatic amines" retrieved from STN Database accession no. 77:19476 XP002233858 abstract & ARCHIV DER PHARMAZIE UND BERICHTE DER DEUTSCHEN PHARMAZEUTISCHEN GESELLSCHAFT (1972), 305(3), 159-71, ----- -/--	1-28, 36-41

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 02/01127

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; YOUNGDALE, GILBERT A. ET AL: "Synthesis and antiinflammatory activity of 5-substituted 2,3-bis(p-methoxyphenyl)indoles" retrieved from STN Database accession no. 71:91199 XP002233859 abstract & JOURNAL OF MEDICINAL CHEMISTRY (1969), 12, 948-9,	1-28, 36-41
P,X	----- WO 02 04425 A (BOEHRINGER INGELHEIM LTD (CA)) 17 January 2002 (2002-01-17) the whole document	1-42
P,X	----- US 2002/065418 A1 (BOEHRINGER INGELHEIM CO (US)) 30 May 2002 (2002-05-30) the whole document	1-42
P,X	----- US 6 358 992 B1 (CELL PATHWAYS INC (US)) 19 March 2002 (2002-03-19) claims; examples -----	1-28, 36-41

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA 02/01127

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 1-25 (part), 28-42 (part)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-25 (part), 28-42 (part)

Present claims 1-25, 28, 29 and 36 relate to an extremely large number of possible compounds. Additionally, claim 1 contain so many provisos that a lack of clarity within the meaning of Article 6 PCT arises to such an extent as to render a meaningful search of the claims impossible. Consequently, the search has been carried out for those parts of the application which do appear to be clear and supported by the examples, namely compounds of the structural features of the compounds in claims 26 and 27, i.e. compounds of formula (I) according to claim 1, wherein:

A is O, S or NR₁; B is CR₃; M is CR₅ or NR₅; K is CR₄; L is CR₇; Y₁ is O; Z is OR₆; R₂ is optionally substituted phenyl or heterocyclyl; R₃ is C₃-C₇ cycloalkyl; the remaining substituents are as defined in claim 1.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 02/01127

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0147883	A	05-07-2001	AU 2401701 A	09-07-2001
			CN 1342145 T	27-03-2002
			CZ 20013424 A3	13-02-2002
			EP 1162196 A1	12-12-2001
			HU 0202263 A2	28-12-2002
			WO 0147883 A1	05-07-2001
			JP 2001247550 A	11-09-2001
			NO 20014134 A	22-10-2001
			NZ 514403 A	25-10-2002
			SK 13752001 A3	02-07-2002
			TR 200103147 T1	21-06-2002
US 5059621	A	22-10-1991	LU 85544 A1	03-04-1986
			US 5428052 A	27-06-1995
			AT 401514 B	25-09-1996
			AT 42192 A	15-02-1996
			AT 395714 B	25-02-1993
			AT 271885 A	15-07-1992
			AU 578310 B2	20-10-1988
			AU 4755985 A	27-03-1986
			AU 593838 B2	22-02-1990
			AU 6299586 A	08-01-1987
			BE 903254 A1	18-03-1986
			BE 904421 A4	30-06-1986
			CA 1256862 A1	04-07-1989
			CA 1256804 A1	04-07-1989
			CH 665841 A5	15-06-1988
			CH 675241 A5	14-09-1990
			DE 3533308 A1	27-03-1986
			DE 3546907 C2	10-06-1998
			DK 169475 B1	07-11-1994
			ES 8608487 A1	01-12-1986
			FR 2570377 A1	21-03-1986
			GB 2164648 A ,B	26-03-1986
			GB 2197320 A ,B	18-05-1988
			GR 852270 A1	21-01-1986
			IT 1209663 B	30-08-1989
			JP 2055068 C	23-05-1996
			JP 7084437 B	13-09-1995
			JP 61085360 A	30-04-1986
			MX 166783 B	04-02-1993
			NL 8502557 A ,B,	16-04-1986
			PT 81132 A ,B	01-10-1985
			SE 461466 B	19-02-1990
			SE 8504291 A	20-03-1986
			SE 502470 C2	23-10-1995
			SE 8800949 A	16-03-1988
			US 4920140 A	24-04-1990
			US 5260295 A	09-11-1993
			US 4740519 A	26-04-1988
			US 5288744 A	22-02-1994
			ZA 8507158 A	28-05-1986
US 6069156	A	30-05-2000	AT 219765 T	15-07-2002
			AU 713460 B2	02-12-1999
			AU 5123496 A	30-10-1996
			CA 2217707 A1	17-10-1996
			DE 69622031 D1	01-08-2002

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 02/01127

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 6069156	A	DE 69622031 T2 EP 0820441 A1 ES 2175079 T3 WO 9632379 A1 JP 11503445 T TW 420663 B ZA 9602859 A	12-12-2002 28-01-1998 16-11-2002 17-10-1996 26-03-1999 01-02-2001 11-10-1996
WO 9748697	A	24-12-1997 AU 3102697 A CA 2258728 A1 EP 0934307 A1 WO 9748697 A1 JP 2000509719 T US 2002173527 A1 US 6303600 B1 ZA 9705446 A	07-01-1998 24-12-1997 11-08-1999 24-12-1997 02-08-2000 21-11-2002 16-10-2001 21-12-1998
JP 10067682	A	10-03-1998 NONE	
JP 10114654	A	06-05-1998 NONE	
JP 09124632	A	13-05-1997 NONE	
WO 0132653	A	10-05-2001 US 6455525 B1 AU 1581101 A CN 1387528 T EP 1226141 A1 NO 20022095 A TR 200201225 T2 WO 0132653 A1	24-09-2002 14-05-2001 25-12-2002 31-07-2002 11-06-2002 21-08-2002 10-05-2001
US 3565912	A	23-02-1971 US 3654308 A	04-04-1972
JP 2001122855	A	08-05-2001 NONE	
WO 0130774	A	03-05-2001 DE 19951360 A1 AU 1272801 A BR 0015026 A CN 1379772 T CZ 20021413 A3 WO 0130774 A1 EP 1261601 A1 NO 20021808 A SK 5432002 A3	03-05-2001 08-05-2001 16-07-2002 13-11-2002 17-07-2002 03-05-2001 04-12-2002 17-04-2002 06-11-2002
US 5912260	A	15-06-1999 AT 197146 T AU 2534295 A AU 682051 B2 AU 7347894 A DE 69426205 D1 DE 69426205 T2 DK 720601 T3 EP 0720601 A1 EP 0763026 A1 ES 2152989 T3 FI 960572 A WO 9504720 A2 WO 9532949 A1	15-11-2000 21-12-1995 18-09-1997 28-02-1995 30-11-2000 08-03-2001 20-11-2000 10-07-1996 19-03-1997 16-02-2001 07-02-1996 16-02-1995 07-12-1995

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 02/01127

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5912260	A	GB 2290539 A	03-01-1996
		GR 3035100 T3	30-03-2001
		HU 75301 A2	28-05-1997
		JP 10504525 T	06-05-1998
		JP 9502430 T	11-03-1997
		NO 960488 A	15-03-1996
		NZ 269827 A	28-10-1996
		PL 312960 A1	27-05-1996
		PT 720601 T	28-02-2001
		SG 52229 A1	28-09-1998
		US 5919829 A	06-07-1999
		US 5795907 A	18-08-1998
		ZA 9504315 A	26-11-1996
US 4740519	A	26-04-1988	
		LU 85544 A1	03-04-1986
		AT 401514 B	25-09-1996
		AT 42192 A	15-02-1996
		AT 395714 B	25-02-1993
		AT 271885 A	15-07-1992
		AU 578310 B2	20-10-1988
		AU 4755985 A	27-03-1986
		AU 593838 B2	22-02-1990
		AU 6299586 A	08-01-1987
		BE 903254 A1	18-03-1986
		BE 904421 A4	30-06-1986
		CA 1256862 A1	04-07-1989
		CA 1256804 A1	04-07-1989
		CH 665841 A5	15-06-1988
		CH 675241 A5	14-09-1990
		DE 3533308 A1	27-03-1986
		DE 3546907 C2	10-06-1998
		DK 169475 B1	07-11-1994
		ES 8608487 A1	01-12-1986
		FR 2570377 A1	21-03-1986
		GB 2164648 A ,B	26-03-1986
		GB 2197320 A ,B	18-05-1988
		GR 852270 A1	21-01-1986
		IT 1209663 B	30-08-1989
		JP 2055068 C	23-05-1996
		JP 7084437 B	13-09-1995
		JP 61085360 A	30-04-1986
		MX 166783 B	04-02-1993
		NL 8502557 A ,B,	16-04-1986
		PT 81132 A ,B	01-10-1985
		SE 461466 B	19-02-1990
		SE 8504291 A	20-03-1986
		SE 502470 C2	23-10-1995
		SE 8800949 A	16-03-1988
		US 5428052 A	27-06-1995
		US 4920140 A	24-04-1990
		US 5059621 A	22-10-1991
		US 5260295 A	09-11-1993
		US 5288744 A	22-02-1994
		ZA 8507158 A	28-05-1986
EP 0987250	A	22-03-2000	
		AU 6229298 A	09-09-1998
		EP 0987250 A1	22-03-2000
		WO 9837069 A1	27-08-1998

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 02/01127

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9928297	A	10-06-1999	DE 19753522 A1	10-06-1999
			AU 2267199 A	16-06-1999
			WO 9928297 A1	10-06-1999
EP 0242167	A	21-10-1987	AT 69812 T	15-12-1991
			AU 602515 B2	18-10-1990
			AU 7139387 A	22-10-1987
			CA 1329933 A1	31-05-1994
			CN 87103504 A	09-12-1987
			DD 258984 A5	10-08-1988
			DE 3774749 D1	09-01-1992
			DK 190587 A	16-10-1987
			EP 0242167 A2	21-10-1987
			ES 2038172 T3	16-07-1993
			FI 871632 A	16-10-1987
			GR 3003209 T3	17-02-1993
			HU 43819 A2	28-12-1987
			IE 60013 B1	18-05-1994
			JP 2035969 C	28-03-1996
			JP 7068211 B	26-07-1995
			JP 63008369 A	14-01-1988
			NO 871588 A	16-10-1987
			NZ 219978 A	28-08-1990
			PL 265151 A1	06-03-1989
			PT 84681 A , B	01-05-1987
			SU 1597098 A3	30-09-1990
			US 4898863 A	06-02-1990
			ZA 8702313 A	25-11-1987
			ZW 6387 A1	23-11-1988
WO 0147922	A	05-07-2001	AU 2209401 A	09-07-2001
			BR 0017038 A	07-01-2003
			CZ 20022207 A3	13-11-2002
			EP 1263759 A2	11-12-2002
			WO 0147922 A2	05-07-2001
			NO 20023032 A	21-06-2002
WO 0204425	A	17-01-2002	AU 7225801 A	21-01-2002
			WO 0204425 A2	17-01-2002
			US 2002065418 A1	30-05-2002
			US 6479508 B1	12-11-2002
			WO 02070739 A2	12-09-2002
US 2002065418	A1	30-05-2002	US 6479508 B1	12-11-2002
			AU 7225801 A	21-01-2002
			WO 0204425 A2	17-01-2002
			WO 02070739 A2	12-09-2002
US 6358992	B1	19-03-2002	NONE	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.